

Interference in the radiation of two point-like charges

Yurij Yaremko*

*Institute for Condensed Matter Physics,
1 Svientsitskii St., 79011 Lviv, Ukraine*

Abstract

Energy-momentum and angular momentum carried by electromagnetic field of two point-like charged particles arbitrarily moving in flat spacetime are presented. Apart from usual contributions to the Noether quantities produced separately by particles 1 and 2, the conservation laws contain also joint contribution due to the fields of both particles. The mixed part of Maxwell energy-momentum density is decomposed into bound and radiative components which are separately conserved off the world lines of particles. The former describes the deformation of electromagnetic clouds of “bare” charges due to mutual interaction while the latter defines the radiation which escapes to infinity. The bound terms contribute to particles’ individual 4-momenta while the radiative ones exert the radiation reaction. Analysis of energy-momentum and angular momentum balance equations results the Lorentz-Dirac equation as an equation of motion for a pointed charge under the influence of its own electromagnetic field as well as field produced by another charge.

PACS numbers: 03.50.De, 11.10.Gh, 11.30.Cp

1 Introduction

The most natural and widely accepted equation of motion for a charge when radiation reaction is taken into account is the Lorentz-Dirac equation [1]. This equation has been discussed mainly for the case of one charge in an external electromagnetic field. In the present paper we consider an isolated system of two point electric charges and their electromagnetic field. We study the electromagnetic energy-momentum and angular momentum radiated by charges; the study of energy-momentum and angular momentum balance equations implies the Lorentz-Dirac equation for more than one charge.

*Electronic mail: yar@ph.icmp.lviv.ua

The dynamics of the entire system is governed by the action

$$I = \sum_{a=1}^2 \left(-m_a \int ds_a \sqrt{-\dot{z}_a^2} + e_a \int ds_a A_{a,\mu} \dot{z}_a^\mu \right) - \frac{1}{16\pi} \int d^4y f_{\mu\nu} f^{\mu\nu}, \quad (1.1)$$

where $f_{\mu\nu} = \sum_a (\partial_\mu A_{a,\nu} - \partial_\nu A_{a,\mu})$ is the total field generated by two charges. Charge e_a moves on a world line $\zeta_a \in \mathbb{M}_4$ described by functions $z_a^\mu(s_a)$ which give the particle's coordinates as functions of proper time s_a ; $\dot{z}_a^\mu := dz_a^\mu/ds_a$ is the a -th four-velocity.

The action is invariant under the space-time translations and rotations which constitute the Poincaré group. This immediately implies conserved quantities which place stringent requirements on the dynamics of the system. They demand that the change in electromagnetic field momentum [2],

$$p_{\text{em}}^\nu = \int_\Sigma d\sigma_\alpha T^{\alpha\nu}, \quad (1.2)$$

and angular momentum [2],

$$M_{\text{em}}^{\mu\nu} = \int_\Sigma d\sigma_\alpha (y^\mu T^{\alpha\nu} - y^\nu T^{\alpha\mu}), \quad (1.3)$$

should be balanced by a corresponding change in the total momentum and total angular momentum of the particles. (By $d\sigma_\alpha$ we denote the vectorial surface element on a space-like hypersurface Σ .)

Since the Maxwell energy-momentum tensor density,

$$4\pi T^{\mu\nu} = f^{\mu\lambda} f^\nu{}_\lambda - 1/4 \eta^{\mu\nu} f^{\kappa\lambda} f_{\kappa\lambda}, \quad (1.4)$$

is quadratic in the field and this field satisfies the superposition principle, the total electromagnetic field stress-energy tensor is

$$T^{\mu\nu} = T_{(1)}^{\mu\nu} + T_{(2)}^{\mu\nu} + T_{\text{int}}^{\mu\nu}, \quad (1.5)$$

where a -th particle density $T_{(a)}^{\mu\nu}$ is given by the expression (1.4) where “total” field strengths $f^{\mu\nu}$ are substituted by “individual” ones $f_{(a)}^{\mu\nu}$. The mixed term

$$4\pi T_{\text{int}}^{\mu\nu} = f_{(1)}^{\mu\lambda} f_{(2)\lambda}^\nu + f_{(2)}^{\mu\lambda} f_{(1)\lambda}^\nu - 1/4 \eta^{\mu\nu} \left(f_{(1)}^{\kappa\lambda} f_{\kappa\lambda}^{(2)} + f_{(2)}^{\kappa\lambda} f_{\kappa\lambda}^{(1)} \right) \quad (1.6)$$

describes the joint contribution due to both fields.

In this paper we study radiation produced by an isolated system of two point electric charges and their electromagnetic field. Outgoing electromagnetic waves remove energy, momentum, and angular momentum from the sources which then undergo radiation reaction. The verification of conservation laws is not a trivial matter, since the interference contribution (1.8) involves divergent terms.

In the derivation of particle's equation of motion, Dirac [1] evaluated the flux of electromagnetic energy-momentum over a narrow world tube surrounding the particle's

world line. The author substituted the components of the retarded Liénard-Wiechert field in the stress-energy tensor (1.4) inside the narrow tube. In 1970 Teitelboim [3] splits the “retarded” stress-energy tensor into the “bound” and “emitted” parts which are separately conserved off the world line of the particle. The author calculates the flow of energy-momentum out of the portion of Bhabha world tube [4] bounded by tilted spacelike hypersurfaces which are orthogonal to particle’s four-velocity at instants τ and $\tau + d\tau$, respectively. Bound part, $\hat{T}_{\text{bnd}}$, describes a rigid electromagnetic “cloud” which are permanently attached to the source and carried along with it. “Bare” charge and “cloud” constitute new entity: dressed charged particle. $\hat{T}_{\text{bnd}}$ contributes into particle’s inertia: 4-momentum of dressed charge contains, apart from usual velocity term, also a term which is proportional to the square of charge e_a ,

$$p_a^\mu = m_a u_a^\mu - \frac{2e_a^2}{3} a_a^\mu. \quad (1.7)$$

(Coulomb-like infinity stemming from the pointness of “bare” source is absorbed by the rest mass m_a within the renormalization procedure.) Time derivative of the second term in eq.(1.7) is the well-known Schott term which describes a reversible form of emission and absorption of field energy, which never gets far from the point-like source. The radiative part, $\hat{T}_{\text{rad}}$, yields the Larmor relativistic rate of radiated energy-momentum. It detaches itself from the charge and leads an independent existence. This rate together with the Schott term constitutes the Abraham radiation reaction vector. López and Villarroel [5] split the torque of the stress-energy tensor into bound and emitted components which possess analogous properties.

The results can be applied to the first and the second terms in the total electromagnetic field stress-energy tensor (1.5) which describe “individual” radiation contributions due to particles 1 and 2, respectively. The question is what part of the mixed density should be taken instead of (1.6) to describe the radiation which reaches to a very distant sphere?

Aguirregabiria and Bel [6] studied the interference part of energy-momentum,

$$p_{\text{int}}^\nu = \int_\Sigma d\sigma_\mu T_{\text{int}}^{\mu\nu}, \quad (1.8)$$

carried by electromagnetic field of two point charges. The authors prove the fundamental theorem that the mixed radiation rate does not depend on the shape of spacelike surface Σ which is used to integrate the mixed part (1.6) of the Maxwell energy-momentum tensor density. For a prescribed plane motion of the charges the perturbation scheme is elaborated within the framework of predictive relativistic mechanics [7, 8]. The lowest approximation gives the well-known expression [9, p.214] for the dipole radiation of two point charges moving according to Coulomb’s law. In Ref. [10] the scheme is applied to the angular momentum carried by electromagnetic field of two point charges.

In the case of N particles we would merely obtain as an obvious generalization of eq. (1.5) the sum of N one-particle terms and the mixed contributions corresponding to $N(N-1)/2$ pairs of charges. Hence, the interference component dominates in the radiation from a bunch of identical charged particles (e.g., in free electron lasers). To evaluate the radiation of a relativistic N -body system, Klepikov [11] defines the center of a system of

radiation events which allows to synchronize the instants at which electromagnetic waves emitted by different charges combine on a very distant sphere. Fourier analysis is applied to calculate the time and angular distributions of energy-momentum flux. The radiation of a bunch of charged particles moving in a uniform magnetic field is considered in detail.

The only exact solution is obtained by Rivera and Villarroel Ref. [12, eq.(3.27)]. The authors calculate the rate of radiation (including interference part) generated by two identical point charges rotating uniformly at opposite ends of a diameter, in a fixed circle. External fields which govern the strictly prescribed motions are constructed. The rate of radiation is evaluated via the retarded Liénard-Wiechert fields produced by the charges. Further [13] more general case of circular motion of two unlike charges in two coplanar and concentric circumferences is considered.

Note that nine years before the radiation by a system of two uniformly circling charges has been evaluated by Hnizdo [14] (see also discussion [15, 16]). The author concludes that “the power radiated by such a system equals exactly the rate at which work is done on the system by external force”.

In this paper we study the interference part of energy-momentum and angular momentum of electromagnetic field generated by two arbitrarily moving charges¹. We restrict ourselves to the retarded Liénard-Wiechert solutions; the advanced ones are rejected on the grounds of causality. In Section 2 we introduce coordinate system which makes relatively easy the calculation of the covariant 4-momentum radiated by interacting charges. The calculation is performed in Sections 3 and 4. We reveal divergence-free radiative part of the mixed density (1.6). It determines the radiation that escapes to infinity while the (short-range) bound part modifies individual 4-momenta (1.7) of dressed particles. The mixed part of radiated energy-momentum depends only on velocities, accelerations and the relative 4-position of the charges. In Section 5 we derive equations of motion of interacting charged particles. Analysis of energy-momentum and angular momentum balance equations results the well-known Lorentz-Dirac equation. In Section 6 we study symmetry properties of radiative energy-momentum and angular momentum which rely on invariance of action (1.1) under inversions of space and time axes. In Section 7 we discuss the results and implications.

2 “Interference” coordinate system

To perform the surface integration (1.8) of interference stress-energy tensor, an appropriate coordinate system is necessary. Such a coordinate system is introduced in Ref. [6]. It involves the evolution parameter λ associated with an inertial observer; the surface of integration is a surface of constant λ . In Refs. [17] and [18] this coordinate system is adapted to the simplest hyperplane $\Sigma_t = \{y \in \mathbb{M}_4 : y^0 = t\}$ associated with an unmoving inertial observer. The “laboratory” time t is a single common parameter defined along all the world lines of the system.

¹The generalization of this work to N charges is an obvious one.




Figure 1: The interference picture in a plane Σ_t . Sphere S_a with radius $k_a^0 = t - t_a$ is centered at point O_a with coordinates $z_a^{i'}(t_a), i' = 1, 2, 3$. Angle φ distinguishes the points of intersection $S_1 \cap S_2 = C(O, h)$ which constitute support of integrals (2.1) and (2.2). k_a^0, k_a^3 , and h are the components of the future oriented null 4-vector $k_a^{\alpha'} = \Omega^{\alpha'}_{\alpha}(y^{\alpha} - z_a^{\alpha}(t_a))$. Matrix $\hat{\Omega}$ determines transition to “momentarily rotating” Lorentz frame.

The mixed contributions to energy-momentum,

$$p_{\text{int}}^{\nu}(t) = \int_{\Sigma_t} d\sigma_0 T_{\text{int}}^{0\nu}, \quad (2.1)$$

and angular momentum,

$$M_{\text{int}}^{\mu\nu}(t) = \int_{\Sigma_t} d\sigma_0 (y^{\mu} T_{\text{int}}^{0\nu} - y^{\nu} T_{\text{int}}^{0\mu}), \quad (2.2)$$

are due to interference of spherical wave fronts S_1 and S_2 in Σ_t (see Fig. 1). Sphere,

$$S_a(\mathbf{z}_a(t_a), t - t_a) = \{y \in \mathbb{M}_4 : (y^0 - t_a)^2 = \sum_i (y^i - z_a^i(t_a))^2, y^0 = t, t - t_a > 0\}, \quad (2.3)$$

is the intersection of the future light cone with vertex at point $z_a(t_a) \in \zeta_a$ and Σ_t . This contribution is zero if the relative position 4-vector $q = z_1 - z_2$ is timelike. If q is spacelike, the intersection $S_1 \cap S_2$ is the circle $C(O, h)$ with radius h ; in “momentarily rotating” Lorentz frame the circle $C(O, h)$ lies in Oxy plane and centered at the coordinate origin (see Fig. 1). If points z_1 and z_2 are related by a null ray, the intersection $S_1 \cap S_2$ contains the only point.

2.1 Local map

To find the local expressions for coordinate transformation $(y^{\alpha}) \mapsto (t, t_1, t_2, \varphi)$, we translate the origin of the laboratory Lorentz frame at the center O of the circle $C(O, h) =$




Figure 2: The past light cone with vertex at point $y \in \Sigma_t$ is punctured by the world lines of the 1-st particle and the 2-nd particle at points $z_1(t_1)$ and $z_2(t_2)$, respectively. The vector K_a is a null vector pointing from the emission point $z_a(t_a) = (t_a, z_a^i(t_a))$ to a field point y . The relative position 4-vector $q = z_1 - z_2$ is equal to difference $K_2 - K_1$; its square $(q \cdot q) = -2(K_2 \cdot K_1)$.

$S_1 \cap S_2$ and then rotate space axes till a new z -axis be directed along 3-vector $\mathbf{q} := \mathbf{z}_1 - \mathbf{z}_2$,

$$y^\alpha = z_a^\alpha(t_a) + \Omega^\alpha{}_{\alpha'}(t_1, t_2)k_a^{\alpha'}. \quad (2.4)$$

Here $k_a, a = 1, 2$ is the future oriented null-vector with components

$$k_a^0 = t - t_a, \quad k_a^1 = h \sin \varphi, \quad k_a^2 = h \cos \varphi, \quad k_a^3 = (-1)^a \frac{q}{2} + \frac{(k_2^0)^2 - (k_1^0)^2}{2q}, \quad (2.5)$$

which arise from analysis of triangle O_1O_2H pictured in Fig. 1 (we denote $q = |\mathbf{q}|$). Matrix space-time components are $\Omega_{0\mu} = \Omega_{\mu 0} = \delta_{\mu 0}$. Its space components Ω_{ij} constitute an orthogonal 3×3 matrix (A.6) which determines the rotation described above (see Appendix A). It defines new orthonormal basis,

$$\begin{aligned} \mathbf{n}_\vartheta &= \cos \varphi_q \cos \vartheta_q \mathbf{e}_1 + \sin \varphi_q \cos \vartheta_q \mathbf{e}_2 - \sin \vartheta_q \mathbf{e}_3, \\ \mathbf{n}_\varphi &= -\sin \varphi_q \mathbf{e}_1 + \cos \varphi_q \mathbf{e}_2, \\ \mathbf{n}_q &= \cos \varphi_q \sin \vartheta_q \mathbf{e}_1 + \sin \varphi_q \sin \vartheta_q \mathbf{e}_2 + \cos \vartheta_q \mathbf{e}_3, \end{aligned} \quad (2.6)$$

which is constructed from components of the relative position 3-vector $\mathbf{q}$, e.g. $\cos \varphi_q = q^1 / \sqrt{(q^1)^2 + (q^2)^2}$, $\cos \vartheta_q = q^3 / q$.

To find the Jacobian of coordinate transformation (2.4), we derive its differential chart. Setting $\alpha = 0$ in eq.(2.4) immediately follows $y^0 = t$. Since $t = y^0$, then $\partial t / \partial y^\alpha = \delta_{0\alpha}$.

Because y and $z_a(t_a)$ lie on the light cone (see Fig. 2), a change field point y comes with a change in t_a . Suppose that y is displaced to the new point $y + \delta y$. The new intersection of the past light cone of this vertex with the a -th world line is then $z_a(t_a + \delta t_a)$. These points are still related by the equation

$$(y^0 + \delta y^0 - t_a - \delta t_a)^2 = \sum_i (y^i + \delta y^i - z_a^i(t_a + \delta t_a))^2. \quad (2.7)$$

Expanding this to the first order in δy and δt_a and using the cone equation (2.3), we obtain $K_{a,\alpha} \delta y^\alpha - (v_a \cdot K_a) \delta t_a = 0$ or

$$\begin{aligned} \frac{\partial t_a}{\partial y^\alpha} &= -\frac{K_{a,\alpha}}{r_a} \\ &= -\frac{\Omega_{\alpha\alpha'} k_a^{\alpha'}}{r_a}. \end{aligned} \quad (2.8)$$

Here K_a is a -th null vector pictured in figure 2 and symbol r_a denotes the scalar product $(v_a \cdot K_a)$, taken with opposite sign; noncovariant 4-velocity $v_a := (1, dz_a^i/dt_a)$.

For the angular variable we have

$$\frac{\partial \varphi}{\partial y^\alpha} = \Omega_{\alpha\alpha'} k_\varphi^{\alpha'}, \quad (2.9)$$

where

$$k_\varphi^0 = 0, \quad k_\varphi^1 = \frac{\cos \varphi}{h}, \quad k_\varphi^2 = -\frac{\sin \varphi}{h}, \quad k_\varphi^3 = 0. \quad (2.10)$$

Recall that h is the radius of the circle $C(O, h) = S_1 \cap S_2$ pictured in Fig. 1.

Determinant of the matrix which defines this differential chart gives the inverse Jacobian: $J^{-1} = q/(r_1 r_2)$. The “interference” surface element,

$$d\sigma_0 = \frac{r_1 r_2}{q} dt_1 dt_2 d\varphi, \quad (2.11)$$

is ill defined if and only if the particles are very close to each other.

2.2 Global mapping

Setting $a = 1$ in eq.(2.4) we obtain the coordinate system centered on an accelerated world line of the first particle. The flat spacetime $\mathbb{M}_4$ is a disjoint union of hyperplanes $\Sigma_t = \{y \in \mathbb{M}_4 : y^0 = t\}$. An interference hyperplane Σ_t is a disjoint union of retarded spheres $S_1(O_1, t - t_1)$ centered at points $O_1 \in \Sigma_t$ with coordinates $\mathbf{z}_1(t_1)$. A sphere is covered by its intersections with spherical wave fronts of the second source. Each circle $S_1 \cap S_2$ can be labelled by the individual time t_2 of the second particle and each point on a given circle can be labelled by its polar angle φ .

Going along the world line ζ_1 we arrive unavoidably at the point $t_1^{ret}(t)$, such that the future light cone of $z_1[t_1^{ret}(t)]$ touches the 2-nd world line at point $z_2(t) \in \Sigma_t$ (see Fig. 4). Light cone of upper vertices do not intersect the second world line at all.

In context with the principle of retarded causality, Σ_t is divided into two regions where outgoing waves sourced by charged particles combine in quite different manner.




Figure 3: For a given t_1 the wave front S_1 is covered by circles $S_1 \cap S_2$ if the parameter t_2 increases from $t_2^{ret}(t_1)$ to $t_2^{adv}(t_1)$. Minimal value labels the vertex of forward light cone which is punctured by ζ_1 at $z_1(t_1)$. The largest sphere S_2^{ret} touches the sphere S_1 at point N . World line ζ_2 punctures the future light cone of $z_1(t_1)$ at point $z_2(t_2^{adv})$. Intersection of S_1 and the smallest sphere S_2^{adv} contains the only point S .

- (i) *Causal*, which is filled up by spheres S_1 of radii larger than or equal to $t - t_1^{ret}(t)$.
- (ii) *Acausal*, where parameter t_1 increases from $t_1^{ret}(t)$ to the instant of observation t . It is the ball bounded by sphere of radius $t - t_1^{ret}(t)$ centered at point $\mathbf{z}_1[t_1^{ret}(t)]$.

2.2.1 Causal region

Causal region is spanned by curvilinear coordinates (2.4) where t_1 increases from $-\infty$ to the instant $t_1^{ret}(t)$. To cover the sphere S_1 where t_1 is fixed we change the parameter t_2 which labels point $z_2(t_2) \in \zeta_2$. The starting point is the solution $t_2^{ret}(t_1)$ of algebraic equation $q^0 = q$ or

$$t_1 - t_2 = q(t_1, t_2), \quad (2.12)$$

where points $z_1(t_1) \in \zeta_1$ and $z_2(t_2^{ret}) \in \zeta_2$ are linked by a null ray. The largest sphere $S_2(O_2^{ret}, t - t_2^{ret})$ touches a given sphere $S_1(O_1, t - t_1)$ at only point N (see Fig. 3). If parameter t_2 increases to $t_2^{adv}(t_1)$ being the solution of algebraic equation $q^0 = -q$ or

$$t_2 - t_1 = q(t_1, t_2), \quad (2.13)$$

the intersection $S_1 \cap S_2^{adv}$ contains the only point S . If parameter t_2 changes from $t_2^{ret}(t_1)$ to $t_2^{adv}(t_1)$, the sphere S_1 is covered by circles $C(O, h) = S_1 \cap S_2$.




Figure 4: The instant $t_1^{\text{ret}}(t)$ is the solution of algebraic equation (2.13) with $t_2 = t$. Sphere $S_1(\mathbf{z}_1^{\text{ret}}, t - t_1^{\text{ret}}(t))$ bounds acausal region. For a given $t_1 \in [t_1^{\text{ret}}(t), t]$ the sphere S_1 with radius $t - t_1$ is a disjoint union of circles $S_1 \cap S_2$ if parameter t_2 increases from $t_2^{\text{ret}}(t_1)$ to $t_2'(t, t_1)$. The maximal value satisfies algebraic equation $k_2^0 + k_1^0 = |\mathbf{q}|$. The sphere S_2' with radius $t - t_2'$ touches S_1 at point S . Intersection $S_1 \cap S_2^{\text{ret}}$ contains the only point N .

2.2.2 Acausal region

Acausal region of an interference hyperplane Σ_t corresponds to the fragments of the world lines which are not related to each other. (By this we mean that the radiation emitted by the first particle during the interval $[t_1^{\text{ret}}(t), t]$ does not come to the second one and vice versa.) Nevertheless, the outgoing waves of these portions of world lines combine in Σ_t .

Acausal region is filled up by spheres $S_1(O_1, t - t_1) \in \Sigma_t$, where $t_1 \in [t_1^{\text{ret}}(t), t]$. The sphere S_1 with fixed t_1 is the disjoint union of circles $C(O, h) = S_1 \cap S_2$ if the parameter t_2 increases from $t_2^{\text{ret}}(t_1)$ to $t_2'(t, t_1)$. The starting point of this interval is still the solution of eq.(2.12) while the maximal value of t_2 satisfies the algebraic equation $k_2^0 + k_1^0 = q$ or

$$2t - t_1 - t_2 = q(t_1, t_2). \quad (2.14)$$

A given sphere $S_1(O_1, t - t_1)$ touches $S_2(O_2', t - t_2')$ at only point S (see Fig. 4).

2.2.3 Surface integration

In an analogous way we construct the coordinate system centered on the world line of the second particle. If $t_2 \in]-\infty, t_2^{\text{ret}}(t)]$ then $t_1 \in [t_1^{\text{ret}}(t_2), t_1^{\text{adv}}(t_2)]$; if $t_2 \in [t_2^{\text{ret}}(t), t]$ then $t_1 \in [t_1^{\text{ret}}(t_2), t_1'(t, t_2)]$, $\varphi \in [0, 2\pi[$. The ends of intervals are defined implicitly by algebraic equations (2.12), (2.13), and (2.14).

The surface integration (2.1) and (2.2) can be performed via the coordinate system

centered on a world line either of the first particle,

$$\left[\int_{-\infty}^{t_1^{ret}(t)} dt_1 \int_{t_2^{ret}(t_1)}^{t_2^{adv}(t_1)} dt_2 + \int_{t_1^{ret}(t)}^t dt_1 \int_{t_2^{ret}(t_1)}^{t'_2(t, t_1)} dt_2 \right] \int_0^{2\pi} d\varphi \frac{r_1 r_2}{q}, \quad (2.15)$$

or of the second particle,

$$\left[\int_{-\infty}^{t_2^{ret}(t)} dt_2 \int_{t_1^{ret}(t_2)}^{t_1^{adv}(t_2)} dt_1 + \int_{t_2^{ret}(t)}^t dt_2 \int_{t_1^{ret}(t_2)}^{t'_1(t, t_2)} dt_1 \right] \int_0^{2\pi} d\varphi \frac{r_1 r_2}{q}. \quad (2.16)$$

To calculate the flows (2.1) of the mixed electromagnetic field energy and momentum which flow across the hyperplane Σ_t , we should integrate the Maxwell energy-momentum tensor density (1.6) over angular variable φ and over time variables t_1 and t_2 .

3 Angular integration of energy-momentum and angular momentum tensor densities

In this Section we trace a series of stages in integration of the mixed Maxwell energy-momentum tensor density over φ . In Appendix A we derive some useful expressions.

In terms of Minkowski coordinates (y^α) the electromagnetic field generated by a -th particle is given by

$$\hat{f}_{(a)} = \frac{e_a}{r_a^2} u_a \wedge \mathbf{k}_a + \frac{e_a}{r_a} [a_a \wedge \mathbf{k}_a + (\mathbf{k}_a \cdot a_a) u_a \wedge \mathbf{k}_a], \quad (3.1)$$

where symbol $\wedge$ denotes the wedge product. We use **sans-serif** symbols for the retarded distance²,

$$\mathbf{r}_a = -\eta_{\alpha\beta} (y^\alpha - z_a^\alpha(s_a)) u_a^\beta(s_a), \quad (3.2)$$

and for the null vector $K_a = y - z_a(s_a)$ rescaled by a factor r_a^{-1} ,

$$\mathbf{k}_a^\alpha = \frac{1}{r_a} (y^\alpha - z_a^\alpha(s_a)). \quad (3.3)$$

To express field strengths in terms of curvilinear coordinates (t, t_1, t_2, φ) , it is advantageous to replace the retarded proper time $s_a(y)$ by evolution parameter t_a . The components of particles' 4-velocities u_a and 4-accelerations a_a , $a = 1, 2$, become [2]

$$u_a^\mu = \gamma_a v_a^\mu(t_a), \quad a_a^\mu = \gamma_a^4 (v_a \cdot \dot{v}_a) v_a^\mu + \gamma_a^2 \dot{v}_a^\mu, \quad (3.4)$$

where 4-vectors $v_a^\mu = (1, v_a^i(t_a))$, $\dot{v}_a^\mu = (0, \dot{v}_a^i(t_a))$ and factor $\gamma_a := [1 - \mathbf{v}_a^2]^{-1/2}$. Substituting these into eq.(3.1) and using the relation $\mathbf{k}_a^\mu = K_a^\mu / r_a$ yields

$$\hat{f}_{(a)} = e_a \left(\frac{v_a \wedge K_a}{r_a^3} c_a + \frac{\dot{v}_a \wedge K_a}{r_a^2} \right), \quad (3.5)$$

²Because the speed of light is set to unity, r_a is equal to the spatial distance between $z_a[s_a^{ret}(y)]$ and y as measured in momentarily comoving Lorentz frame where $u_a^\alpha = (1, 0, 0, 0)$.

where

$$r_a = K_a^0 - (\mathbf{K}_a \mathbf{v}_a), \quad c_a = \gamma_a^{-2} + (\mathbf{K}_a \dot{\mathbf{v}}_a). \quad (3.6)$$

Note that r_a is the retarded distance (3.2) rescaled by a factor γ_a , i.e. $r_a = \gamma_a^{-1} \mathbf{r}_a$. The separation vector K_a has the form $\hat{\Omega} k_a$, where components of null vector k_a are given by eqs.(2.5) and matrix $\hat{\Omega}$ determines the transition to momentarily comoving Lorentz frame associated with basis (2.6).

It is straightforward to substitute the components of electromagnetic fields (3.5) in terms of “interference” coordinates (t, t_1, t_2, φ) into integrands of expressions (2.1) and (2.2) to calculate the interference part of radiated energy-momentum and angular momentum, respectively. Integration of the mixed stress-energy tensor over angular variable is the key to the problem. All φ -dependent terms are concentrated in the following constructions:

$$\begin{aligned} \mathcal{D}^a &= \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{a}{q r_1 r_2}, & \mathcal{B}^a &= \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{a c_2}{q r_1 (r_2)^2}, \\ \mathcal{C}^a &= \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{a c_1}{q (r_1)^2 r_2}, & \mathcal{A}^a &= \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{a c_1 c_2}{q (r_1)^2 (r_2)^2}. \end{aligned} \quad (3.7)$$

They are labeled according to their dependence on the combination of components of the separation vectors K_1 and K_2 : factor a is replaced by $K_1^\mu K_2^\nu$, K_1^μ , K_2^ν , or 1 for $\mathcal{D}_{12}^{\mu\nu}$, $\mathcal{D}_1^\mu$, $\mathcal{D}_2^\nu$ or $\mathcal{D}^0$, respectively. (The others $\mathcal{B}^a$, $\mathcal{C}^a$, and $\mathcal{A}^a$ are marked analogously.)

The mixed part of the stress-energy tensor (1.6) is symmetric in indices 1 and 2. Substituting (3.5) into the first term of this expression and using the identities $K_2 - K_1 = q$ and $(K_2 \cdot K_1) = -1/2(q \cdot q)$ yields

$$\begin{aligned} \frac{1}{4\pi} \int_0^{2\pi} d\varphi \ J f_{(1)}^{\mu\alpha} f_{(2)\alpha}^\nu &= \frac{e_1 e_2}{2} \left\{ \mathcal{T}_{12}^{\mu\nu} \left(\frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) + \mathcal{T}_1^\mu \left(v_2^\nu \frac{\partial \sigma}{\partial t_1} \right) + \mathcal{T}_2^\nu \left(v_1^\mu \frac{\partial \sigma}{\partial t_2} \right) + \mathcal{T}^0 (v_1^\mu v_2^\nu \sigma) \right. \\ &- \mathcal{C}_1^\mu v_2^\nu \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} - \mathcal{D}_1^\mu v_2^\nu \frac{\partial^3 \sigma}{\partial t_1^2 \partial t_2} - \mathcal{B}_2^\nu v_1^\mu \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} - \mathcal{D}_2^\nu v_1^\mu \frac{\partial^3 \sigma}{\partial t_1 \partial t_2^2} \\ &- \left. \mathcal{B}^0 v_1^\mu v_2^\nu \frac{\partial \sigma}{\partial t_1} - \mathcal{C}^0 v_1^\mu v_2^\nu \frac{\partial \sigma}{\partial t_2} - \mathcal{D}^0 \left(v_1^\mu v_2^\nu \frac{\partial \sigma}{\partial t_2} + v_1^\mu \dot{v}_2^\nu \frac{\partial \sigma}{\partial t_1} + v_1^\mu v_2^\nu \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) \right\}, \end{aligned} \quad (3.8)$$

after addition of similar terms and integration over φ . World function $\sigma(t_1, t_2)$ of two spacelike related points, $z_1(t_1) \in \zeta_1$ and $z_2(t_2) \in \zeta_2$, is equal to one-half of the square of vector $q = z_1 - z_2$, taken with opposite sign,

$$\sigma(t_1, t_2) = -1/2(q \cdot q). \quad (3.9)$$

Each second order differential operator,

$$\hat{T}^a = \mathcal{D}^a \frac{\partial^2}{\partial t_1 \partial t_2} + \mathcal{B}^a \frac{\partial}{\partial t_1} + \mathcal{C}^a \frac{\partial}{\partial t_2} + \mathcal{A}^a, \quad (3.10)$$

has been labeled according to dependence of coefficients (3.7) on the combination of vectors K_1 and K_2 .

For the convolution $f_{(1)}^{\alpha\beta} f_{\alpha\beta}^{(2)}$, we obtain

$$\frac{1}{4\pi} \int_0^{2\pi} d\varphi J f_{(1)}^{\alpha\beta} f_{\alpha\beta}^{(2)} = e_1 e_2 \mathcal{T}^0(\lambda), \quad (3.11)$$

where function

$$\lambda = \sigma \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} - \frac{\partial \sigma}{\partial t_1} \frac{\partial \sigma}{\partial t_2} \quad (3.12)$$

depends on two-point function (3.9) and its derivatives in time variables.

To distinguish the partial derivatives in time variables, we rewrite the operator (3.10) as the sum of the second-order differential operator,

$$\hat{\Pi}^a = \frac{\partial^2}{\partial t_1 \partial t_2} \mathcal{D}^a + \frac{\partial}{\partial t_1} \left(\mathcal{B}^a - \frac{\partial \mathcal{D}^a}{\partial t_2} \right) + \frac{\partial}{\partial t_2} \left(\mathcal{C}^a - \frac{\partial \mathcal{D}^a}{\partial t_1} \right), \quad (3.13)$$

and the “tail”,

$$\pi^a = \frac{\partial^2 \mathcal{D}^a}{\partial t_1 \partial t_2} - \frac{\partial \mathcal{B}^a}{\partial t_1} - \frac{\partial \mathcal{C}^a}{\partial t_2} + \mathcal{A}^a. \quad (3.14)$$

For a smooth function $f(t_1, t_2)$ we have

$$\hat{\mathcal{T}}^a(f) = \hat{\Pi}^a(f) + f \pi^a. \quad (3.15)$$

Cumbersome calculations which are presented in Appendix A give the relations

$$\begin{aligned} \pi^0 &= 0, \\ \pi_1^\mu &= v_1^\mu \left(\mathcal{B}^0 - \frac{\partial \mathcal{D}^0}{\partial t_2} \right), \quad \pi_2^\nu = v_2^\nu \left(\mathcal{C}^0 - \frac{\partial \mathcal{D}^0}{\partial t_1} \right), \\ \pi_{12}^{\mu\nu} &= v_1^\mu \left(\mathcal{B}_2^\nu - \frac{\partial \mathcal{D}_2^\nu}{\partial t_2} \right) + v_2^\nu \left(\mathcal{C}_1^\mu - \frac{\partial \mathcal{D}_1^\mu}{\partial t_1} \right) - v_1^\mu v_2^\nu \mathcal{D}^0, \end{aligned} \quad (3.16)$$

which allow us to rewrite the sum of integrals (3.8) and (3.11) in terms of differential operators $\hat{\Pi}^a$ and partial derivatives in t_1 and t_2 ,

$$\begin{aligned} \mathcal{P}_{12}^{\mu\nu} &= \frac{1}{4\pi} \int_0^{2\pi} d\varphi J \left(f_{(1)}^{\mu\alpha} f_{(2)\alpha}^\nu - \frac{\eta^{\mu\nu}}{4} f_{(1)}^{\alpha\beta} f_{\alpha\beta}^{(2)} \right) \\ &= \frac{e_1 e_2}{2} \left\{ \hat{\Pi}_{12}^{\mu\nu} \left(\frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) + \hat{\Pi}_1^\mu \left(v_2^\nu \frac{\partial \sigma}{\partial t_1} \right) + \hat{\Pi}_2^\nu \left(v_1^\mu \frac{\partial \sigma}{\partial t_2} \right) + \hat{\Pi}^0(v_1^\mu v_2^\nu \sigma) \right. \\ &\quad - \frac{\partial}{\partial t_1} \left(\mathcal{D}_1^\mu v_2^\nu \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) - \frac{\partial}{\partial t_2} \left(\mathcal{D}_2^\nu v_1^\mu \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) \\ &\quad \left. - \frac{\partial}{\partial t_1} \left(\mathcal{D}^0 v_1^\mu v_2^\nu \frac{\partial \sigma}{\partial t_2} \right) - \frac{\partial}{\partial t_2} \left(\mathcal{D}^0 v_1^\mu v_2^\nu \frac{\partial \sigma}{\partial t_1} \right) - \frac{1}{2} \eta^{\mu\nu} \hat{\Pi}^0(\lambda) \right\}. \end{aligned} \quad (3.17)$$

We denote $\mathcal{P}_{21}^{\mu\nu}$ the integral over φ of the remaining terms involved in tensor (1.6). It can be obtained by interchanging of indices 1 and 2.

Setting $\mu = 0$ and $\nu = i$ in eq.(3.17), we obtain the first term of the mixed space-time components of the stress-energy tensor (1.6). We add the term where indices 1 and 2 are interchanged. Since zeroth components k_1^0 and k_2^0 of the separation four-vectors K_1 and K_2 do not depend on φ , the final expression get simplified,

$$\begin{aligned}\mathcal{P}_{\text{int}}^i &= \mathcal{P}_{12}^{0i} + \mathcal{P}_{21}^{0i} \\ &= \frac{e_1 e_2}{2} \left[\hat{\Pi}_2^i \left(\frac{\partial \lambda_1}{\partial t_2} \right) + \hat{\Pi}_1^i \left(\frac{\partial \lambda_2}{\partial t_1} \right) + \hat{\Pi}^0 (v_2^i \lambda_1 + v_1^i \lambda_2) \right. \\ &\quad \left. - \frac{\partial}{\partial t_1} \left(v_2^i \frac{\partial \lambda_1}{\partial t_2} \mathcal{D}^0 \right) - \frac{\partial}{\partial t_2} \left(v_1^i \frac{\partial \lambda_2}{\partial t_1} \mathcal{D}^0 \right) \right],\end{aligned}\tag{3.18}$$

where

$$\lambda_1 = k_1^0 \frac{\partial \sigma}{\partial t_1} + \sigma, \quad \lambda_2 = k_2^0 \frac{\partial \sigma}{\partial t_2} + \sigma.\tag{3.19}$$

Similarly we derive zeroth component $\mathcal{P}_{\text{int}}^0$. Setting $\mu = 0$ and $\nu = 0$ in eq.(3.17), we obtain the first one-half of desired expression. The second one, $\mathcal{P}_{21}^{00}$, can be derived via interchanging indices 1 and 2. The integral of energy density T_{int}^{00} over the angular variable has the form

$$\begin{aligned}\mathcal{P}_{\text{int}}^0 &= \mathcal{P}_{12}^{00} + \mathcal{P}_{21}^{00} \\ &= \frac{e_1 e_2}{2} \hat{\Pi}^0 \left(\Sigma \frac{\partial^2 \Sigma}{\partial t_1 \partial t_2} - \frac{\partial \Sigma}{\partial t_1} \frac{\partial \Sigma}{\partial t_2} \right),\end{aligned}\tag{3.20}$$

where three-point function,

$$\Sigma(t, t_1, t_2) = 2k_1^0 k_2^0 + \sigma(t_1, t_2),\tag{3.21}$$

depends on particles' positions referred to the moments t_1 and t_2 before observation instant t as well as on t itself.

We now turn to the integration of the angular momentum tensor density (2.2) carried by the electromagnetic field due to two pointlike charges. We present the torque $m_{\text{int}}^{\mu\nu} = y^\mu T_{\text{int}}^{0\nu} - y^\nu T_{\text{int}}^{0\mu}$ in the following form:

$$m_{\text{int}}^{\mu\nu} = m_{12}^{\mu\nu} + m_{21}^{\mu\nu} - m_{12}^{\nu\mu} - m_{21}^{\nu\mu},\tag{3.22}$$

where

$$m_{12}^{\mu\nu} = (z_1^\mu + K_1^\mu) \frac{1}{4\pi} \left[f_{(1)}^{0\lambda} f_{(2)\lambda}^\nu - \frac{1}{4} \eta^{0\nu} f_{(1)}^{\alpha\beta} f_{\alpha\beta}^{(2)} \right].\tag{3.23}$$

It is straightforward to substitute the fields (3.5) into this expression to calculate the first term of expression (3.22). The others can be obtained by interchanging of the pair of indices (1, 2) and (μ, ν) .

Having integrated expression $Jm_{12}^{\mu\nu}$ over φ we obtain

$$\begin{aligned}
\mathcal{M}_{12}^{\mu\nu} = & \frac{e_1 e_2}{2} \left\{ \hat{T}_{12}^{\mu\nu} \left(\frac{\partial \lambda_1}{\partial t_2} \right) + \hat{T}_1^\mu (v_2^\nu \lambda_1) - v_2^\nu \frac{\partial \lambda_1}{\partial t_2} \mathcal{C}_1^\mu - v_2^\nu \frac{\partial^2 \lambda_1}{\partial t_1 \partial t_2} \mathcal{D}_1^\mu \right. \\
& + \hat{T}_2^\nu \left(z_1^\mu \frac{\partial \lambda_1}{\partial t_2} \right) - v_1^\mu \frac{\partial \lambda_1}{\partial t_2} \mathcal{B}_2^\nu - v_1^\mu \frac{\partial^2 \lambda_1}{\partial t_2^2} \mathcal{D}_2^\nu + \hat{T}^0 (z_1^\mu v_2^\nu \lambda_1) - v_1^\mu v_2^\nu \lambda_1 \mathcal{B}^0 \\
& - v_1^\mu \left(v_2^\nu \lambda_1 + v_2^\nu \frac{\partial \lambda_1}{\partial t_2} \mathcal{D}^0 \right) - z_1^\mu v_2^\nu \frac{\partial \lambda_1}{\partial t_2} \mathcal{C}^0 - z_1^\mu v_2^\nu \frac{\partial^2 \lambda_1}{\partial t_1 \partial t_2} \mathcal{D}^0 \\
& \left. - \frac{\eta^{0\nu}}{2} \left[\hat{T}_1^\mu (\lambda) + \hat{T}^0 (z_1^\mu \lambda) - v_1^\mu \lambda \mathcal{B}^0 - v_1^\mu \frac{\partial \lambda}{\partial t_2} \mathcal{D}^0 \right] \right\}, \tag{3.24}
\end{aligned}$$

where functions λ and λ_a are given by eqs.(3.12) and (3.19), respectively.

Usage of the equalities in eq.(3.16) derived in Appendix A allows us to rewrite the integrand (3.24) as follows:

$$\begin{aligned}
\mathcal{M}_{12}^{\mu\nu} = & \frac{e_1 e_2}{2} \left\{ \hat{\Pi}_{12}^{\mu\nu} \left(\frac{\partial \lambda_1}{\partial t_2} \right) + \hat{\Pi}_1^\mu (v_2^\nu \lambda_1) - \frac{\partial}{\partial t_1} \left(v_2^\nu \frac{\partial \lambda_1}{\partial t_2} \mathcal{D}_1^\mu \right) \right. \\
& + \hat{\Pi}_2^\nu \left(z_1^\mu \frac{\partial \lambda_1}{\partial t_2} \right) - \frac{\partial}{\partial t_2} \left(v_1^\mu \frac{\partial \lambda_1}{\partial t_2} \mathcal{D}_2^\nu \right) \\
& + \hat{\Pi}^0 (z_1^\mu v_2^\nu \lambda_1) - \frac{\partial}{\partial t_1} \left(z_1^\mu v_2^\nu \frac{\partial \lambda_1}{\partial t_2} \mathcal{D}^0 \right) - \frac{\partial}{\partial t_2} (v_1^\mu v_2^\nu \lambda_1 \mathcal{D}^0) \\
& \left. - \frac{\eta^{0\nu}}{2} \left[\hat{\Pi}_1^\mu (\lambda) + \hat{\Pi}^0 (z_1^\mu \lambda) - \frac{\partial}{\partial t_2} (v_1^\mu \lambda \mathcal{D}^0) \right] \right\}. \tag{3.25}
\end{aligned}$$

The other terms of mixed angular momentum,

$$\mathcal{M}_{\text{int}}^{\mu\nu} = \mathcal{M}_{12}^{\mu\nu} + \mathcal{M}_{21}^{\mu\nu} - \mathcal{M}_{12}^{\nu\mu} - \mathcal{M}_{21}^{\nu\mu}, \tag{3.26}$$

can be obtained via interchanging of indices (1, 2) and (μ, ν) .

We see that the integration of the mixed stress-energy tensor (1.6) over φ yields the combinations of partial derivatives in time variables. In the next Section we classify them and reveal the long-range terms which contribute into radiated energy-momentum.

4 Radiative parts of mixed energy-momentum and angular momentum

In previous Section we integrate the mixed part of the stress-energy tensor and its torque over polar angle. Resulted expressions describe contributions to electromagnetic field's energy-momentum and angular momentum due to interference of spherical wave fronts of charges e_1 and e_2 placed at fixed points $z_1(t_1) \in \zeta_1$ and $z_2(t_2) \in \zeta_2$, respectively. The crucial issue is that the integrals (3.18), (3.20) and (3.26) have the remarkable property of being the sum of partial derivatives in time variables. This circumstance allows us to calculate how much electromagnetic field's energy-momentum and angular momentum flow across a hyperplane Σ_t .

It is natural to integrate the expression being the time derivative with respect to t_2 according to the rule (2.15),

$$\begin{aligned}\mathcal{G}_2 &= \left[\int_{-\infty}^{t_1^{ret}(t)} dt_1 \int_{t_2^{ret}(t_1)}^{t_2^{adv}(t_1)} dt_2 + \int_{t_1^{ret}(t)}^t dt_1 \int_{t_2^{ret}(t_1)}^{t'_2(t, t_1)} dt_2 \right] \frac{\partial G_2(t_1, t_2)}{\partial t_2} \\ &= \int_{-\infty}^{t_1^{ret}(t)} dt_1 G_2[t_1, t_2^{adv}(t_1)] - \int_{-\infty}^t dt_1 G_2[t_1, t_2^{ret}(t_1)] + \int_{t_1^{ret}(t)}^t dt_1 G_2[t_1, t'_2(t, t_1)].\end{aligned}\quad (4.1)$$

Having applied the rule (2.16) to the expression of type $\partial G_1/\partial t_1$, we obtain

$$\begin{aligned}\mathcal{G}_1 &= \left[\int_{-\infty}^{t_2^{ret}(t)} dt_2 \int_{t_1^{ret}(t_2)}^{t_1^{adv}(t_2)} dt_1 + \int_{t_2^{ret}(t)}^t dt_2 \int_{t_1^{ret}(t_2)}^{t'_1(t, t_2)} dt_1 \right] \frac{\partial G_1(t_1, t_2)}{\partial t_1} \\ &= \int_{-\infty}^{t_2^{ret}(t)} dt_2 G_1[t_1^{adv}(t_2), t_2] - \int_{-\infty}^t dt_2 G_1[t_1^{ret}(t_2), t_2] + \int_{t_2^{ret}(t)}^t dt_2 G_1[t'_1(t, t_2), t_2].\end{aligned}\quad (4.2)$$

The end points are valuable only in the integration procedure. They are solutions of algebraic equations (2.12), (2.13) and (2.14). The retarded instants $t_a^{ret}(t_b)$ and advanced ones $t_b^{adv}(t_a)$ label the points N and S in which fronts of *outgoing* electromagnetic waves produced by charges touch each other (see Fig. 3). All the moments are *before* the observation instant t , so that the retarded causality is not violated.

It is worth noting that the functions $t_1^{ret}(t_2)$ and $t_2^{adv}(t_1)$ are inverted to each other as well as the pair of functions $t_1^{adv}(t_2)$ and $t_2^{ret}(t_1)$. For a fixed laboratory time t the functions $t'_1(t, t_2)$ and $t'_2(t, t_1)$ are inverses too. These circumstances allow us to change the variables $t_a \mapsto t_a^{ret}(t_b)$ in the “advanced” integrals in eqs.(4.1) and (4.2). Further we couple them with their “retarded” counterparts. Since

$$\frac{dt_1^{ret}(t_2)}{dt_2} = \frac{1 - (\mathbf{v}_1 \mathbf{n}_q)}{1 - (\mathbf{v}_2 \mathbf{n}_q)}, \quad \frac{dt_2^{ret}(t_1)}{dt_1} = \frac{1 + (\mathbf{v}_2 \mathbf{n}_q)}{1 + (\mathbf{v}_1 \mathbf{n}_q)}, \quad (4.3)$$

we obtain

$$\begin{aligned}\mathcal{G}_2 + \mathcal{G}_1 &= \int_{-\infty}^t dt_1 \left[\frac{1 - (\mathbf{v}_1 \mathbf{n}_q)}{1 - (\mathbf{v}_2 \mathbf{n}_q)} G_1 - G_2 \right]_{t_2=t_2^{ret}(t_1)} \\ &\quad + \int_{-\infty}^t dt_2 \left[-G_1 + \frac{1 + (\mathbf{v}_2 \mathbf{n}_q)}{1 + (\mathbf{v}_1 \mathbf{n}_q)} G_2 \right]_{t_1=t_1^{ret}(t_2)} \\ &\quad + \int_{t_2^{ret}(t)}^t dt_2 \left[G_1 + \frac{1 - (\mathbf{v}_2 \mathbf{n}_q)}{1 + (\mathbf{v}_1 \mathbf{n}_q)} G_2 \right]_{t_1=t'_1(t, t_2)}.\end{aligned}\quad (4.4)$$

The unit vector $\mathbf{n}_q = \mathbf{q}/q$ is the third vector of orthonormal triad (2.6). The last integral is due to interference of outgoing electromagnetic waves radiated out by the particles within the acausal region (see Fig. 4). We take into account that

$$\frac{dt'_2(t, t_1)}{dt_1} = -\frac{1 + (\mathbf{v}_1 \mathbf{n}_q)}{1 - (\mathbf{v}_2 \mathbf{n}_q)}. \quad (4.5)$$

Of course, one can change the variables $t_b \mapsto t_b^{adv}(t_a)$ in the “retarded” integrals in eqs.(4.1) and (4.2) and add them to their “advanced” counterparts,

$$\begin{aligned} \mathcal{G}_2 + \mathcal{G}_1 &= \int_{-\infty}^{t_2^{ret}(t)} dt_2 \left[G_1 - \frac{1 - (\mathbf{v}_2 \mathbf{n}_q)}{1 - (\mathbf{v}_1 \mathbf{n}_q)} G_2 \right]_{t_1=t_1^{adv}(t_2)} \\ &+ \int_{-\infty}^{t_1^{ret}(t)} dt_1 \left[-\frac{1 + (\mathbf{v}_1 \mathbf{n}_q)}{1 + (\mathbf{v}_2 \mathbf{n}_q)} G_1 + G_2 \right]_{t_2=t_2^{adv}(t_1)} \\ &+ \int_{t_1^{ret}(t)}^t dt_1 \left[\frac{1 + (\mathbf{v}_1 \mathbf{n}_q)}{1 - (\mathbf{v}_2 \mathbf{n}_q)} G_1 + G_2 \right]_{t_2=t'_2(t, t_1)}. \end{aligned} \quad (4.6)$$

A combination of the “retarded” and the “advanced” terms is valuable too.

Integral of a mixed double derivative can be written in the form either

$$\begin{aligned} \mathcal{I}_1 &= \left[\int_{-\infty}^{t_2^{ret}(t)} dt_2 \int_{t_1^{ret}(t_2)}^{t_1^{adv}(t_2)} dt_1 + \int_{t_2^{ret}(t)}^t dt_2 \int_{t_1^{ret}(t_2)}^{t'_1(t, t_2)} dt_1 \right] \frac{\partial}{\partial t_1} \left[\frac{\partial G(t_1, t_2)}{\partial t_2} \right] \\ &= \int_{-\infty}^{t_2^{ret}(t)} dt_2 \left[\frac{\partial G(t_1, t_2)}{\partial t_2} \right]_{t_1=t_1^{adv}(t_2)} - \int_{-\infty}^t dt_2 \left[\frac{\partial G(t_1, t_2)}{\partial t_2} \right]_{t_1=t_1^{ret}(t_2)} \\ &+ \int_{t_2^{ret}(t)}^t dt_2 \left[\frac{\partial G(t_1, t_2)}{\partial t_2} \right]_{t_1=t'_1(t, t_2)}, \end{aligned} \quad (4.7)$$

or

$$\begin{aligned} \mathcal{I}_2 &= \left[\int_{-\infty}^{t_1^{ret}(t)} dt_1 \int_{t_2^{ret}(t_1)}^{t_2^{adv}(t_1)} dt_2 + \int_{t_1^{ret}(t)}^t dt_1 \int_{t_2^{ret}(t_1)}^{t'_2(t, t_1)} dt_2 \right] \frac{\partial}{\partial t_2} \left[\frac{\partial G(t_1, t_2)}{\partial t_1} \right] \\ &= \int_{-\infty}^{t_1^{ret}(t)} dt_1 \left[\frac{\partial G(t_1, t_2)}{\partial t_1} \right]_{t_2=t_2^{adv}(t_1)} - \int_{-\infty}^t dt_1 \left[\frac{\partial G(t_1, t_2)}{\partial t_1} \right]_{t_2=t_2^{ret}(t_1)} \\ &+ \int_{t_1^{ret}(t)}^t dt_1 \left[\frac{\partial G(t_1, t_2)}{\partial t_1} \right]_{t_2=t'_2(t, t_1)}. \end{aligned} \quad (4.8)$$




Figure 5: Function $G[t_1, t_2^{ret}(t_1)]$ is associated with the interference of wave fronts $S_1(O_1, t - t_1)$ and $S_2(O_2^{ret}, t - t_2^{ret})$ at point $N \in \Sigma_t$. Function $G[t_1, t_2'(t, t_1)]$ is connected with the combination of wave fronts $S_1(O_1, t - t_1)$ and $S_2(O_2', t - t_2')$ at point $S \in \Sigma_t$. If $t_1 \rightarrow t$, both the point N and the point S tend to $z_1(t) = \zeta_1 \cap \Sigma_t$. If $\lim_{t_1 \rightarrow t} G[t_1, t_2^{ret}(t_1)] = \lim_{t_1 \rightarrow t} G[t_1, t_2'(t, t_1)]$ the value of integral of mixed derivative $\partial^2 G / \partial t_1 \partial t_2$ over time variables does not depend on the order of integration.

The question is what expression should be used.

To compare $\mathcal{I}_1$ and $\mathcal{I}_2$ we change the variables in the “advanced” integrals and subtract (4.8) and (4.7). We arrive at the integrals being functions of the end points only,

$$\begin{aligned} \mathcal{I}_1 - \mathcal{I}_2 &= G[t_1, t_2^{ret}(t_1)] \Big|_{t_1 \rightarrow -\infty}^{t_1 \rightarrow t} - G[t_1^{ret}(t_2), t_2] \Big|_{t_2 \rightarrow -\infty}^{t_2 \rightarrow t} \\ &+ \begin{cases} \text{either} & G[t_1'(t, t_2), t_2] \Big|_{t_2 \rightarrow t_2^{ret}(t)}^{t_2 \rightarrow t} \\ \text{or} & - G[t_1, t_2'(t, t_1)] \Big|_{t_1 \rightarrow t_1^{ret}(t)}^{t_1 \rightarrow t} \end{cases}. \end{aligned} \quad (4.9)$$

It vanishes if and only if (i) limiting values of G (that evaluated at the remote past) cancel each other, and (ii) function G is smooth at points at which the world lines puncture Σ_t . Indeed, the part of difference (4.9) which depends on the momentary state of particles’ motion can be rewritten as follows:

$$\begin{aligned} \mathcal{I}_1 - \mathcal{I}_2 &= \lim_{t_1 \rightarrow t} \{ G[t_1, t_2^{ret}(t_1)] - G[t_1, t_2'(t, t_1)] \} \\ &- \lim_{t_2 \rightarrow t} \{ G[t_1^{ret}(t_2), t_2] - G[t_1'(t, t_2), t_2] \}. \end{aligned} \quad (4.10)$$

The situation is illustrated in Fig. 5.

4.1 Criteria

The main task of the present paper is to decompose the interference part of the Maxwell energy-momentum tensor density into bound and radiative components. The former modifies individual 4-momenta (1.7) of dressed particles while the latter shows how a charge is influenced by radiation of another charge.

To reveal meaningful radiative part of mixed energy-momentum (2.1) and angular momentum (2.2), we apply the criteria which were first formulated in Ref. [3, Table 1].

- The bound part diverges while the radiative one is finite.
- The bound component depends on the momentary state of the particles' motion while the radiative one is accumulated with time.
- The form of the bound terms heavily depends on choosing of an integration surface while the radiative terms are invariant.

There are, however, a several properties that the desired expressions have possess before they can be accepted. We list them below.

1. Radiative parts should be completely determined by particles' motion; they can not depend on distance to a point of observation.
2. They should be produced by divergence-free expressions.
3. Non-accelerated charges do not radiate.
4. Balance of Noether conserved quantities yields the Lorentz-Dirac equation.

4.2 Radiative part of mixed momentum

To decompose the momentum (3.18) into bound and radiative components is a straight-forward integration of all the terms over time variables. Scrupulous computations reveal candidate for radiative part,

$$\begin{aligned} \mathcal{P}_{\text{int,rad}}^i &= \frac{e_1 e_2}{2} \left[\hat{\Pi}_2^i \left(k_1^0 \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) - \frac{\partial}{\partial t_1} \left(v_2^i k_1^0 \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \mathcal{D}^0 \right) \right. \\ &\quad \left. + \hat{\Pi}_1^i \left(k_2^0 \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) - \frac{\partial}{\partial t_2} \left(v_1^i k_2^0 \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \mathcal{D}^0 \right) \right], \end{aligned} \quad (4.11)$$

which produce the terms that satisfy Teitelboim's criteria.

Equations (4.1) and (4.2) imply that its integral over time variables is completely determined by values of the arguments of time differential operators at the ends of integration intervals. When a consideration is restricted to the end points where radius h of intersection $S_1 \cap S_2$ vanishes, the coefficients (3.7) get simplified. Since eq. (2.5) the

integrands do not depend on the polar angle at all. The integral over φ becomes unit operator. Radiative momentum (4.11) contains also non-trivial constructions,

$$\mathcal{C}^a - \frac{\partial \mathcal{D}^a}{\partial t_1}, \quad \mathcal{B}^a - \frac{\partial \mathcal{D}^a}{\partial t_2}, \quad (4.12)$$

which are calculated in Appendix A. All the expressions (A.35), (A.41), (A.43), and (A.53) are regular if $h = 0$. Values of relative position 4-vector q , retarded distances r_a , and some basic functions at limiting points are collected in Table 1 (see Appendix B).

The sum of differential operators $\hat{\Pi}_1^i$ and $\hat{\Pi}_2^i$ contains mixed double derivative of the function,

$$G^i = k_1^0 \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \mathcal{D}_2^i + k_2^0 \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \mathcal{D}_1^i. \quad (4.13)$$

A surprising feature of integration of expression (4.11) over time variables is that the result heavily depends on the order of differentiation in $\partial^2 G^i / \partial t_1 \partial t_2$. If one choose the rule (4.7) they obtain

$$p_{\text{rad},21}^i = \int_{-\infty}^t dt_1 \gamma_1^{-1} G_{21}^i[t_1, t_2^{\text{ret}}(t_1)] + \int_{-\infty}^{t_1^{\text{ret}}(t)} dt_1 \gamma_1^{-1} G_{21}^i[t_1, t_2^{\text{adv}}(t_1)]. \quad (4.14)$$

Having applied the rule (4.8) we arrive at

$$p_{\text{rad},12}^i = \int_{-\infty}^t dt_2 \gamma_2^{-1} G_{12}^i[t_1^{\text{ret}}(t_2), t_2] + \int_{-\infty}^{t_2^{\text{ret}}(t)} dt_2 \gamma_2^{-1} G_{12}^i[t_1^{\text{adv}}(t_2), t_2]. \quad (4.15)$$

Function under integral signs,

$$\begin{aligned} \gamma_a^{-1} G_{ba}^i &= e_a e_b \left[\frac{q_{ab}^i(v_a \cdot v_b)}{r_b^3} \gamma_b^{-2} + \frac{q_{ab}^i(v_a \cdot v_b)}{r_b^3} (q_{ab} \cdot \dot{v}_b) \right. \\ &\quad \left. + \frac{q_{ab}^i(v_a \cdot \dot{v}_b)}{r_b^2} - \frac{v_b^i(v_a \cdot v_b)}{r_b^2} \right], \end{aligned} \quad (4.16)$$

is referred to the retarded and the advanced instants. Relative distance 4-vector $q_{ab} := z_a - z_b$; in our notation $q_{12} = q$ while $q_{21} = -q$. Denominator $r_b = -(q_{ab} \cdot v_b)$ is the retarded distance of type (3.6) where both the field point and the point of emission are placed on particles' world lines. So, in expression (4.14) the field point is $z_1(t_1) \in \zeta_1$ where e_1 is placed. Points of emission are $z_2[t_2^{\text{ret}}(t_1)] \in \zeta_2$ in the first path integral and $z_2[t_2^{\text{adv}}(t_1)] \in \zeta_2$ in the second one,

$$r_2^{\text{ret}} = q[1 - (\mathbf{v}_2 \mathbf{n}_q)] \Big|_{[t_1, t_2^{\text{ret}}(t_1)]}, \quad r_2^{\text{adv}} = -q[1 + (\mathbf{v}_2 \mathbf{n}_q)] \Big|_{[t_1, t_2^{\text{adv}}(t_1)]}. \quad (4.17)$$

In expression (4.15) the field point is in $z_2(t_2) \in \zeta_2$ where the second charge is placed. It is acted on by the first charge placed at points either $z_1[t_1^{\text{ret}}(t_2)] \in \zeta_1$ or $z_1[t_1^{\text{adv}}(t_2)] \in \zeta_1$. The distances are as follows:

$$r_1^{\text{ret}} = q[1 + (\mathbf{v}_1 \mathbf{n}_q)] \Big|_{[t_1^{\text{ret}}(t_2), t_2]}, \quad r_1^{\text{adv}} = -q[1 - (\mathbf{v}_1 \mathbf{n}_q)] \Big|_{[t_1^{\text{adv}}(t_2), t_2]}. \quad (4.18)$$

It is of great importance that the integrand (4.16) does not depend on the distances to points N or S on the observation hyperplane Σ_t . It depends on the relative distance between charged particles, on their velocities, and on their accelerations. The situation looks like a -th charge is acted on by b -th one directly. The charges are connected by a null ray: in the first integral in eq.(4.14) or in eq.(4.15) the interaction be forward while in the second one backward in time. All the moments are before the instant of observation t which labels the upper limits of path integrals.

Since the function (4.13) is not smooth in neighborhoods of intersections $\zeta_a \cap \Sigma_t$, the expressions (4.14) and eq.(4.15) are not equivalent. By virtue of the relation (4.9) we compute the difference:

$$p_{\text{rad},21}^i - p_{\text{rad},12}^i = - q^i (A_{21}^{\text{ret}} \cdot A_{12}^{\text{adv}}) \Big|_{[t_1, t_2^{\text{ret}}(t_1)]}^{t_1=t} - q^i (A_{12}^{\text{ret}} \cdot A_{21}^{\text{adv}}) \Big|_{[t_1^{\text{ret}}(t_2), t_2]}^{t_2=t} \Big|_{t_2 \rightarrow -\infty}^{t_2=t} . \quad (4.19)$$

Symbolically we denote

$$A_{ba}^{\text{ret}} = e_b \frac{v_b[t_b^{\text{ret}}(t_a)]}{r_b[t_a, t_b^{\text{ret}}(t_a)]}, \quad A_{ba}^{\text{adv}} = e_b \frac{v_b[t_b^{\text{adv}}(t_a)]}{r_b[t_a, t_b^{\text{adv}}(t_a)]} \quad (4.20)$$

the Liénard-Wiechert vector potential of b -th charge at point at which a -th one is placed.

4.3 Radiative part of mixed stress-energy tensor

In this subsection we find the terms which produce the radiative part of mixed momentum, either (4.14) or (4.15). We start with φ -momentum (4.11) which is then nothing but the mixed space-time component of the following tensor:

$$\begin{aligned} \mathcal{P}_{\text{int},\text{rad}}^{\mu\nu} &= \frac{e_1 e_2}{2} \left[\hat{\Pi}_{12}^{\mu\nu} \left(\frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) - \frac{\partial}{\partial t_1} \left(\mathcal{D}_1^\mu v_2^\nu \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) - \frac{\partial}{\partial t_2} \left(\mathcal{D}_2^\nu v_1^\mu \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) \right. \\ &\quad \left. + \hat{\Pi}_{21}^{\mu\nu} \left(\frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) - \frac{\partial}{\partial t_2} \left(\mathcal{D}_2^\mu v_1^\nu \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) - \frac{\partial}{\partial t_1} \left(\mathcal{D}_1^\nu v_2^\mu \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) \right]. \quad (4.21) \end{aligned}$$

To restore φ -dependent terms leading to this expression, we insert

$$\hat{\Pi}_{ab}^{\mu\nu}(f) = \hat{\mathcal{T}}_{ab}^{\mu\nu}(f) - f \pi_{ab}^{\mu\nu} \quad (4.22)$$

and substitute the right hand side of the third line of eqs.(3.16) for $\pi_{ab}^{\mu\nu}$. (Operator $\hat{\mathcal{T}}_{ab}^{\mu\nu}$ is defined by eq.(3.10).) After cancellation of like terms we obtain a linear combination of coefficients (3.7). Further we omit the integration over polar angle and multiply the result on inverse Jacobian $J^{-1} = q/(r_1 r_2)$. Finally, we obtain the tensor,

$$\begin{aligned} 4\pi t^{\mu\nu} &= e_1 e_2 \left\{ (K_1^\mu K_2^\nu + K_2^\mu K_1^\nu) \left[\frac{c_1 c_2}{r_1^3 r_2^3} \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} + \frac{c_1}{r_1^3 r_2^2} \frac{\partial^3 \sigma}{\partial t_1 \partial t_2^2} + \frac{c_2}{r_1^2 r_2^3} \frac{\partial^3 \sigma}{\partial t_1^2 \partial t_2} + \frac{1}{r_1^2 r_2^2} \frac{\partial^4 \sigma}{\partial t_1^2 \partial t_2^2} \right] \right. \\ &\quad - (v_1^\mu K_2^\nu + K_2^\mu v_1^\nu) \left[\frac{c_2}{r_1^2 r_2^3} \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} + \frac{1}{r_1^2 r_2^2} \frac{\partial^3 \sigma}{\partial t_1 \partial t_2^2} \right] \\ &\quad - (K_1^\mu v_2^\nu + v_2^\mu K_1^\nu) \left[\frac{c_1}{r_1^3 r_2^2} \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} + \frac{1}{r_1^2 r_2^2} \frac{\partial^3 \sigma}{\partial t_1^2 \partial t_2} \right] \\ &\quad \left. + (v_1^\mu v_2^\nu + v_2^\mu v_1^\nu) \frac{1}{r_1^2 r_2^2} \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right\}, \quad (4.23) \end{aligned}$$

which is symmetric in its indices.

Similarly to the mixed stress-energy tensor (1.6) itself, its radiative part,

$$T_{\text{int,rad}}^{\mu\nu} = t^{\mu\nu} - \frac{1}{2}\eta^{\mu\nu}t^\alpha{}_\alpha, \quad (4.24)$$

contains the convolution of tensor (4.23),

$$\begin{aligned} 4\pi t^\alpha{}_\alpha = & 2e_1e_2 \left\{ \frac{c_1c_2}{r_1^3r_2^3}\sigma\frac{\partial^2\sigma}{\partial t_1\partial t_2} + \frac{c_1}{r_1^3r_2^2}\frac{\partial}{\partial t_2}\left(\sigma\frac{\partial^2\sigma}{\partial t_1\partial t_2}\right) + \frac{c_2}{r_1^2r_2^3}\frac{\partial}{\partial t_1}\left(\sigma\frac{\partial^2\sigma}{\partial t_1\partial t_2}\right) \right. \\ & + \frac{1}{r_1^2r_2^2}\frac{\partial^2}{\partial t_1\partial t_2}\left(\sigma\frac{\partial^2\sigma}{\partial t_1\partial t_2}\right) + \frac{c_2}{r_1r_2^3}\frac{\partial^2\sigma}{\partial t_1\partial t_2} + \frac{1}{r_1r_2^2}\frac{\partial^3\sigma}{\partial t_1\partial t_2^2} \\ & \left. + \frac{c_1}{r_1^3r_2}\frac{\partial^2\sigma}{\partial t_1\partial t_2} + \frac{1}{r_1^2r_2}\frac{\partial^3\sigma}{\partial t_1^2\partial t_2} \right\}. \end{aligned} \quad (4.25)$$

Integration over φ results in the combination of partial derivatives in time variables,

$$\int_0^{2\pi} d\varphi Jt^\alpha{}_\alpha = e_1e_2 \left[\hat{\Pi}^0 \left(\sigma\frac{\partial^2\sigma}{\partial t_1\partial t_2} \right) + \frac{\partial}{\partial t_1} \left(\frac{1}{q\|\mathbf{r}_1\|} \frac{\partial^2\sigma}{\partial t_1\partial t_2} \right) + \frac{\partial}{\partial t_2} \left(\frac{1}{q\|\mathbf{r}_2\|} \frac{\partial^2\sigma}{\partial t_1\partial t_2} \right) \right], \quad (4.26)$$

where expressions

$$\|\mathbf{r}_a\| = \sqrt{[k_a^0 - (\mathbf{v}_a\mathbf{n}_q)k_a^3]^2 - h^2[\mathbf{v}_a\mathbf{n}_q]^2}, \quad (4.27)$$

$a = 1, 2$, are introduced in Appendix A.

Finally, after a tedious calculations³ we derive the identity $\partial_\nu T_{\text{int,rad}}^{\mu\nu} = 0$. It means that the radiative part (4.24) of mixed energy-momentum tensor density (1.6) is conserved off particles' world lines.

It is worth noting that the terms in the first line of expression (4.23) only belong to the mixed part (1.6) of the electromagnetic field stress-energy tensor. The others provide vanishing of divergence of $\hat{T}_{\text{int,rad}}$ as well as reasonable expression for radiated energy-momentum which escapes to infinity. By means of the relations

$$v_a^\mu = \gamma_a^{-1}u_a^\mu, \quad \dot{v}_a^\mu = \gamma_a^{-2}(a_a^\mu - \gamma_a^{-1}a_a^0u_a^\mu) \quad (4.28)$$

$$\begin{aligned} r_a &= \gamma_a^{-1}\mathbf{r}_a, & K_a^\mu &= \mathbf{r}_a k_a^\mu, \\ c_a &= \gamma_a^{-2} [1 + \mathbf{r}_a(k_a \cdot a_a) + \gamma_a^{-1}a_a^0\mathbf{r}_a], \end{aligned} \quad (4.29)$$

the expression (4.23) can be easily rewritten in manifestly covariant notations.

4.4 Radiative part of mixed energy

Since $\eta^{00} = -1$, the convolution (4.26) contributes into zeroth component of radiative part of mixed energy-momentum,

$$\begin{aligned} \mathcal{P}_{\text{int,rad}}^0 &= e_1e_2 \left\{ \hat{\Pi}^0 \left[\left(k_1^0k_2^0 + \frac{1}{2}\sigma \right) \frac{\partial^2\sigma}{\partial t_1\partial t_2} \right] \right. \\ &\quad \left. + \frac{1}{2}\frac{\partial}{\partial t_1} \left(\frac{1}{q\|\mathbf{r}_1\|} \frac{\partial^2\sigma}{\partial t_1\partial t_2} \right) + \frac{1}{2}\frac{\partial}{\partial t_2} \left(\frac{1}{q\|\mathbf{r}_2\|} \frac{\partial^2\sigma}{\partial t_1\partial t_2} \right) \right\}. \end{aligned} \quad (4.30)$$

³Differentiation of $T_{\text{int,rad}}$ is straightforward: one can apply the rule (2.8) and combine like terms scaled as $r_1^{-m}r_2^{-n}$; exponents m and n run from 1 to 4 and their sum $3 \leq m+n \leq 7$.

As could be expected the argument of mixed double derivative,

$$G^0 = \left(k_1^0 k_2^0 + \frac{1}{2} \sigma \right) \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \mathcal{D}^0, \quad (4.31)$$

is not smooth in neighborhoods of end points $\zeta_a \cap \Sigma_t$,

$$\Delta \mathcal{I}^0 = - q^0 (A_{21}^{ret} \cdot A_{12}^{adv}) \Big|_{[t_1, t_2^{ret}(t_1)]} \Big|_{t_1 \rightarrow -\infty}^{t_1=t} - q^0 (A_{12}^{ret} \cdot A_{21}^{adv}) \Big|_{[t_1^{ret}(t_2), t_2]} \Big|_{t_2 \rightarrow -\infty}^{t_2=t}. \quad (4.32)$$

Hence the interference rate of radiated energy heavily depends on order of differentiation, either $\partial^2 G^0 / \partial t_1 \partial t_2$ or $\partial^2 G^0 / \partial t_2 \partial t_1$. Having used the rule (4.7) we arrive at the expression of type (4.14). Choosing the rule (4.8) we obtain the expression of type (4.15). The results can be generalized as follows:

$$p_{\text{rad},ba}^\alpha = \int_{-\infty}^t dt_a \gamma_a^{-1} G_{ba}^\alpha [t_a, t_b^{ret}(t_a)] + \int_{-\infty}^{t_a^{ret}(t)} dt_a \gamma_a^{-1} G_{ba}^\alpha [t_a, t_b^{adv}(t_a)], \quad (4.33)$$

where

$$\begin{aligned} \gamma_a^{-1} G_{ba}^\alpha &= e_a e_b \left\{ \frac{q_{ab}^\alpha (v_a \cdot v_b)}{r_b^3} \gamma_b^{-2} + \frac{q_{ab}^\alpha (v_a \cdot v_b)}{r_b^3} (q_{ab} \cdot \dot{v}_b) \right. \\ &\quad \left. + \frac{q_{ab}^\alpha (v_a \cdot \dot{v}_b)}{r_b^2} - \frac{v_b^\alpha (v_a \cdot v_b)}{r_b^2} \right\}. \end{aligned} \quad (4.34)$$

It is convenient to rewrite eq.(4.33) in a manifestly covariant fashion,

$$p_{\text{rad},ab}^\alpha = \int_{-\infty}^{\tau_b} ds_b G_{ab}^\alpha [s_a^{ret}(s_b), s_b] + \int_{-\infty}^{\tau_b^{ret}(\tau_a)} ds_b G_{ab}^\alpha [s_a^{adv}(s_b), s_b]. \quad (4.35)$$

The particles' world lines ζ_1 and ζ_2 are parametrized by individual proper times s_1 and s_2 , respectively. Upper limits τ_1 and τ_2 label the points at which ζ_1 and ζ_2 puncture the observation hyperplane Σ_t . Two-point function,

$$G_{ab}^\alpha = e_a e_b \left\{ \frac{q_{ba}^\alpha (u_b \cdot u_a)}{r_a^3} [1 + (q_{ba} \cdot a_a)] + \frac{q_{ba}^\alpha (u_b \cdot a_a)}{r_a^2} - \frac{u_a^\alpha (u_b \cdot u_a)}{r_a^2} \right\}, \quad (4.36)$$

is evaluated at points on the world lines of particle a and particle b which are linked by a null ray. $u_a(s_a)$ denotes the normalized 4-velocity of a -th particle; 4-acceleration $a_a^\alpha = du_a^\alpha / ds_a$.

In contrast with “one-particle” contributions to radiated energy-momentum, the mixed one is not uniquely defined. Having generalized the expressions (4.19) and (4.32), we obtain

$$p_{\text{rad},21}^\alpha - p_{\text{rad},12}^\alpha = Q_{21}^\alpha \Big|_{s_2 \rightarrow -\infty}^{s_2=\tau_2} - Q_{12}^\alpha \Big|_{s_1 \rightarrow -\infty}^{s_1=\tau_1}, \quad (4.37)$$

where

$$Q_{ab}^\alpha = q_{ab}^\alpha (A_{ab}^{adv} \cdot A_{ba}^{ret}) \Big|_{[s_a, s_b^{ret}(s_a)]}. \quad (4.38)$$

This q -directed null vector is proportional to the scalar product of the Liénard-Wiechert potentials (4.20).

The difference (4.37) depends on the state of particles' motion at the end points of path integrals. Let us evaluate the net energy-momentum which escapes to infinity. The integrals over entire world lines should be substituted for the integrals over the past motion. We suppose that the particles are asymptotically free. Since Liénard-Wiechert potentials fall off at large distances inversely as the first power of the separation vector between the charges, the right-hand side of expression (4.37) vanishes. Hence the full amount of radiative energy-momentum emitted by interacting particles does not depend on the method of integration.

4.5 Radiative part of mixed angular momentum

When an one-particle problem is considered, bound components of the stress-energy tensor [3] and its torque [5] contribute into individual particle's 4-momentum and angular momentum, respectively. The corrections arise from the regularisation procedure which involves the Taylor expansion of surface integrals in which the first two terms lead to the diverging Coulomb self-energy and the Abraham radiation reaction 4-vector, respectively. The finite terms depend on the form of the hole that is cut out from the integration hypersurface to ensure regularization. The best suited hole must be coordinate-free one. Teitelboim [3] integrate over hyperplane $\sigma(\tau) = \{y \in \mathbb{M}_4 : u_\mu(\tau)(y^\mu - z^\mu(\tau)) = 0\}$ which is orthogonal to the 4-velocity of the charge at point $z(\tau)$ at which particle's world line punctures $\sigma(\tau)$. It is evident that the tilted hyperplane together with the future light cone cutting out the coordinate-free hole (see Refs. [3, Fig.1], [5, Fig.1], [2, Fig.5-2]). The considerations lead to manifestly covariant and structure-independent finite terms of clear physical sense.

The difficulties associated with the computation of the mixed contribution (1.6) are twofold — to perform the meaningful decomposition of $\hat{T}_{\text{int}}$ into bound and radiative parts and to choose an appropriate surface of integration. The tilted hyperplane which plays privileged role in the one-particle radiation reaction problem is not suitable whenever two-body one is considered. Indeed, there is no a hyperplane which is orthogonal to the world lines of both the particles at all events. Kosyakov [19] constructs a piecewise hypersurface where a small fragment of a spacelike hyperplane Σ is replaced by a fragment of an orthogonal hyperplane $\sigma_a(\tau_a) = \{y \in \mathbb{M}_4 : u_{a,\mu}(\tau_a)(y^\mu - z_a^\mu(\tau_a)) = 0\}$ in the vicinity of every intersection point. The deformed hyperplane is called *locally adjusted*. But the problem arises how to sew these fragments with Σ .

Expressions (B.4), (B.5), (B.7), and (B.8) show that the surface integrals of bound component of the mixed part of stress-energy tensor depend on the state of particles' motion in vicinities of intersection points $\zeta_a \cap \Sigma_t$. Unfortunately, the Taylor expansions of the divergent components of bound energy-momentum do not lead to reasonable finite covariant terms. It is because the integration surface Σ_t is tightly connected with the laboratory inertial frame. This choice yields the coordinate-dependent hole around a -th particle in the point of intersection $\zeta_a \cap \Sigma_t$. For this reason we assume that an intrinsic structure of a charged particle is beyond the limits of classical theory [2]. We do not require any assumptions about the particle structure, its charge distribution and its size (except that its “radius” does not vanish, although it is too small to be observed). To

reconcile the theory with observation, we assume that a dressed charged particle possesses *finite* 4-momentum and angular momentum.

We face a problem of how the 4-momenta of interacting dressed charges depend on their individual characteristics such as their masses, charges, 4-velocities, etc. Valuable information can be extracted from the radiative part of electromagnetic field's angular momentum. Indeed, conserved quantities place stringent requirements on the dynamics of our closed system. They demand that the change in radiated momentum and angular momentum should be balanced by a corresponding change in the individual momenta and angular momenta of dressed particles. It is felt that analysis of balance equations yields reasonable expressions. Indeed, it is shown [20] that the Lorentz-Dirac equation can be derived from the energy-momentum and angular momentum balance equations. In [21] the analog of the Lorentz-Dirac equation in six dimensions is obtained via analysis of 21 conserved quantities which correspond to the Poincaré symmetry of an isolated point particle coupled with electromagnetic field.

Recall from Section 3 that the integral of mixed angular momentum tensor density over φ is the combination of partial derivatives in time variables. To reveal meaningful radiative terms we integrate eq.(3.26) over t_1 and t_2 and apply Teitelboim's criteria.

The calculation is virtually identical to that presented above, and we shall not bother with details. The result depends on the method of integration of mixed double derivatives,

$$\begin{aligned}\hat{M}_{\text{rad},ab} &= \int_{-\infty}^{\tau_b} ds_b (z_b \wedge G_{ab}^{\text{ret}} - e_b u_b \wedge A_{ab}^{\text{ret}}) \\ &+ \int_{-\infty}^{\tau_b^{\text{ret}}(\tau_a)} ds_b (z_b \wedge G_{ab}^{\text{adv}} - e_b u_b \wedge A_{ab}^{\text{adv}}).\end{aligned}\tag{4.39}$$

(Symbol $\wedge$ denotes the wedge product.) If we choose the rule (4.8) we obtain $\hat{M}_{\text{rad},21}$. If one prefer the rule (4.7) they derive $\hat{M}_{\text{rad},12}$. Direct calculations show that the difference between these expressions depends on the momentary state of particles' motion,

$$\hat{M}_{\text{rad},21} - \hat{M}_{\text{rad},12} = z_2 \wedge Q_{21}|_{s_2 \rightarrow -\infty}^{s_2=\tau_2} - z_1 \wedge Q_{12}|_{s_1 \rightarrow -\infty}^{s_1=\tau_1}.\tag{4.40}$$

Expression (4.39) for radiated angular momentum satisfies Teitelboim criteria. Indeed, the integrand is finite and covariant; the radiation is accumulated with time. Non-covariant bound terms which are presented in Appendix B are quite different: they depend on the state of particles' motion at points $\zeta_a \cap \Sigma_t$ and they contain divergences. We assume that the expression (4.39) is involved in the angular momentum balance equation explicitly.

4.6 Radiation of non-accelerated charges

Let us consider a specific case of very massive particles, such that $e_1/m_1 \ll 1$ and $e_2/m_2 \ll 1$. Electromagnetic field is too small to accelerate the charges so that the particles move with constant velocities. We place the coordinate origin of the Lorentz inertial frame at point at which the second charge is placed. If $\mathbf{v}_2 = 0$, the zeroth

component $\gamma_1^{-1}G_{21}^0$ of two-point function (4.34) is identically equal to zero. Hence the radiative energy,

$$p_{\text{rad},21}^0 = \int_{-\infty}^t dt_1 \gamma_1^{-1} G_{21}^0[t_1, t_2^{\text{ret}}(t_1)] + \int_{-\infty}^{t_1^{\text{ret}}(t)} dt_1 \gamma_1^{-1} G_{21}^0[t_1, t_2^{\text{adv}}(t_1)], \quad (4.41)$$

vanishes.

If $\mathbf{v}_2 = 0$, the space components of this function get simplified,

$$\gamma_1^{-1} G_{21}^i = e_1 e_2 \frac{q^i}{q^3}. \quad (4.42)$$

The function evaluated at advanced instant $t_2^{\text{adv}}(t_1)$ is just the function referred to the retarded instant $t_2^{\text{ret}}(t_1)$ taken with opposite sign: $G_{21}^i[t_1, t_2^{\text{adv}}(t_1)] = -G_{21}^i[t_1, t_2^{\text{ret}}(t_1)]$. Because of shift in limits of the retarded and the advanced integrals the mixed part of radiative 3-momentum is as follows:

$$\begin{aligned} p_{\text{rad},21}^i &= \int_{-\infty}^t dt_1 \gamma_1^{-1} G_{21}^i[t_1, t_2^{\text{ret}}(t_1)] + \int_{-\infty}^{t_1^{\text{ret}}(t)} dt_1 \gamma_1^{-1} G_{21}^i[t_1, t_2^{\text{adv}}(t_1)] \\ &= e_1 e_2 \int_{t_1^{\text{ret}}(t)}^t dt_1 \frac{q^i}{q^3}. \end{aligned} \quad (4.43)$$

This is vanishingly small quantity. It should be rejected if for no other reason than that the radiated 4-momentum can not be spacelike 4-vector.

In a like manner we calculate the radiative angular momentum,

$$\begin{aligned} \hat{M}_{\text{rad},21}^{0i} &= e_1 e_2 \int_{t_1^{\text{ret}}(t)}^t dt_1 \left(t_1 \frac{q^i}{q^3} + \frac{v_1^i}{q} \right), \\ \hat{M}_{\text{rad},21}^{ij} &= 0. \end{aligned} \quad (4.44)$$

If the upper limit of integration $t \rightarrow +\infty$, $p_{\text{rad},21}^i = 0$. Indeed, in this case for each point on ζ_1 there are points on ζ_2 labeled by the retarded and the advanced instants. Corresponding contributions cancel each other.

Alternative expression for mixed contribution to radiated energy-momentum,

$$p_{\text{rad},12}^\alpha = \int_{-\infty}^{+\infty} dt_2 \gamma_2^{-1} G_{12}^\alpha[t_1^{\text{ret}}(t_2), t_2] + \int_{-\infty}^{+\infty} dt_2 \gamma_2^{-1} G_{12}^\alpha[t_1^{\text{adv}}(t_2), t_2], \quad (4.45)$$

differs from $p_{\text{rad},21}^\alpha$ on the sum of null vectors,

$$e_1 e_2 \frac{n_q^\alpha}{q[1 - (\mathbf{n}_q \mathbf{v}_1)]} \Big|_{t_1 \rightarrow -\infty}^{t_1 \rightarrow +\infty} + e_1 e_2 \frac{n_q^\alpha}{q[1 + (\mathbf{n}_q \mathbf{v}_1)]} \Big|_{t_2 \rightarrow -\infty}^{t_2 \rightarrow +\infty}, \quad (4.46)$$

estimated at limits of integrals. If the particles move with different velocities, they are asymptotically free. If both the particles are static, the difference is equal to zero because the distance q between charges does not change.

5 Equations of motion of radiating charges

In this Section we study the energy-momentum and angular momentum balance equations. We calculate how much electromagnetic field energy, momentum, and angular momentum flow across hyperplane Σ_t . We can do it at a time $t + \Delta t$. We demand that the change in these quantities be balanced by a corresponding change in those of the particles, so that the total energy-momentum,

$$P = p_1 + p_2 + \frac{2e_1^2}{3} \int_{-\infty}^{\tau_1} ds_1 a_1^2 u_1(s_1) + \frac{2e_2^2}{3} \int_{-\infty}^{\tau_2} ds_2 a_2^2 u_2(s_2) + p_{\text{int,rad}}, \quad (5.1)$$

and total angular momentum,

$$\begin{aligned} \hat{M} &= z_1 \wedge p_1 + z_2 \wedge p_2 \\ &+ \frac{2e_1^2}{3} \int_{-\infty}^{\tau_1} ds_1 (a_1^2 z_1 \wedge u_1 + u_1 \wedge a_1) + \frac{2e_2^2}{3} \int_{-\infty}^{\tau_2} ds_2 (a_2^2 z_2 \wedge u_2 + u_2 \wedge a_2) + \hat{M}_{\text{int,rad}}, \end{aligned} \quad (5.2)$$

are properly conserved.

We suppose that particles' individual 4-momenta p_1 and p_2 are already renormalized. The words “already renormalized” mean that momenta contain contributions due to bound component of the stress-energy tensor density, including its mixed part. While the radiation which detaches itself from charges and leads an independent existence is involved explicitly.

In contrast with “one-particle” contributions to conserved quantities (5.1) and (5.2), the mixed ones, (4.35) and (4.39), are not uniquely defined. Our theory faces in the radiation problem a significant issue: non-uniqueness in determination of radiation sourced by the mixed part of Maxwell energy-momentum tensor density. To solve the problem we mix the obtained expressions

$$p_{\text{int,rad}} = \kappa p_{\text{rad},21} + (1 - \kappa) p_{\text{rad},12}, \quad (5.3)$$

$$\hat{M}_{\text{int,rad}} = \kappa \hat{M}_{\text{rad},21} + (1 - \kappa) \hat{M}_{\text{rad},12}, \quad (5.4)$$

and try to find out the value of constant κ that accords with experience.

Expressions $p_{\text{rad},ab}$ and $\hat{M}_{\text{rad},ab}$ are based on two-point function (4.36) referred to the points on the world lines of particle a and particle b which are linked by a null ray,

$$\begin{aligned} \sigma &= -\frac{1}{2} (z_1 - z_2)^2 \\ &= 0. \end{aligned} \quad (5.5)$$

This implies that a displacement of $z_1(s_1)$ typically induces a simultaneous displacement of $z_2(s_2)$ because new points $z_1(s_1 + \delta s_1)$ and $z_2(s_2 + \delta s_2)$ must also be linked by a null geodesic,

$$(q_{21} \cdot u_1) ds_1 + (q_{12} \cdot u_2) ds_2 = 0. \quad (5.6)$$

This immediately gives

$$\frac{ds_a}{ds_b} = -\frac{(q_{ba} \cdot u_b)}{r_a}, \quad (5.7)$$

where $r_a = -(q_{ba} \cdot u_a)$.

By virtue of this expression we compare the retarded and the advanced integrals involved in eqs. (4.35) and (4.39). With understanding that functions $s_a^{adv}(s_b)$ and $s_b^{ret}(s_a)$ are inverses, after some algebra we obtain

$$\begin{aligned} \int_{-\infty}^{\tau_b^{ret}(\tau_a)} ds_b G_{ab}^\alpha[s_a^{adv}(s_b), s_b] &= \int_{-\infty}^{\tau_a} ds_a G_{ba}^\alpha[s_a, s_b^{ret}(s_a)] + Q_{ab}^\alpha \Big|_{s_a \rightarrow -\infty}^{s_a = \tau_a}, \quad (5.8) \\ \int_{-\infty}^{\tau_b^{ret}(\tau_a)} ds_b (z_b \wedge G_{ab}^{adv} - e_b u_b \wedge A_{ab}^{adv}) &= \int_{-\infty}^{\tau_a} ds_a (z_a \wedge G_{ba}^{ret} - e_a u_a \wedge A_{ba}^{ret}) \\ &+ z_a \wedge Q_{ab} \Big|_{s_a \rightarrow -\infty}^{s_a = \tau_a}. \quad (5.9) \end{aligned}$$

The relations accord with the right-hand sides of eqs. (4.37) and (4.40).

Equipped with these relations we rewrite the mixed parts of radiated energy-momentum (4.35) and angular momentum (4.39) as follows:

$$p_{\text{rad},ab} = \int_{-\infty}^{\tau_b} ds_b G_{ab}[s_a^{ret}(s_b), s_b] + \int_{-\infty}^{\tau_a} ds_a G_{ba}[s_a, s_b^{ret}(s_a)] + Q_{ab} \Big|_{s_a \rightarrow -\infty}^{s_a = \tau_a}, \quad (5.10)$$

$$\begin{aligned} \hat{M}_{\text{rad},ab} &= \int_{-\infty}^{\tau_b} ds_b (z_b \wedge G_{ab}^{ret} - e_b u_b \wedge A_{ab}^{ret}) \\ &+ \int_{-\infty}^{\tau_a} ds_a (z_a \wedge G_{ba}^{ret} - e_a u_a \wedge A_{ba}^{ret}) + z_a \wedge Q_{ab} \Big|_{s_a \rightarrow -\infty}^{s_a = \tau_a}. \quad (5.11) \end{aligned}$$

Since $Q_{ab} \neq Q_{ba}$, the expressions are not symmetric in indices a and b .

Substituting eqs. (5.3) and (5.4) into right-hand sides of eqs. (5.1) and (5.2), respectively, we obtain the total energy-momentum,

$$P = \sum_{b=1}^2 \left\{ p_b + \frac{2e_b^2}{3} \int_{-\infty}^{\tau_b} ds_b a_b^2 u_b + \int_{-\infty}^{\tau_b} ds_b G_{ab}^{ret} + \kappa_b Q_{ba} \Big|_{s_b \rightarrow -\infty}^{s_b = \tau_b} \right\}, \quad (5.12)$$

and angular momentum,

$$\begin{aligned} \hat{M} &= \sum_{b=1}^2 \left\{ z_b \wedge p_b + \frac{2e_b^2}{3} \int_{-\infty}^{\tau_b} ds_b (a_b^2 z_b \wedge u_b + u_b \wedge a_b) \right. \\ &+ \left. \int_{-\infty}^{\tau_b} ds_b (z_b \wedge G_{ab}^{ret} - e_b u_b \wedge A_{ab}^{ret}) + \kappa_b z_b \wedge Q_{ba} \Big|_{s_b \rightarrow -\infty}^{s_b = \tau_b} \right\}. \quad (5.13) \end{aligned}$$

(In our notations $\kappa_1 := 1 - \kappa$ and $\kappa_2 := \kappa$.) The balance equations are differential consequences of these conserved quantities. Since the action is not propagated instantaneously, the balance in a vicinity of the first charge as well as in a neighborhood of the second charge should be achieved separately,

$$\dot{p}_a = -\frac{2e_a^2}{3} a_a^2 u_a - G_{ba}^{ret} - \kappa_a \dot{Q}_{ab} \quad (5.14)$$

$$u_a \wedge \left(p_a + \frac{2e_a^2}{3} a_a - e_a A_{ba}^{ret} + \kappa_a Q_{ab} \right) = 0. \quad (5.15)$$

(The overdot means the derivation with respect to individual proper time τ_a .) Solution of six linear equations (5.15) in four components of a -th 4-momentum contains an arbitrary scalar function, say m_a ,

$$p_a = m_a u_a - \frac{2e_a^2}{3} a_a + e_a A_{ba}^{\text{ret}} - \kappa_a Q_{ab}. \quad (5.16)$$

Since $(u_a \cdot a_a) = 0$, the scalar product of momentum of a -th particle on its 4-acceleration is as follows:

$$(p_a \cdot a_a) = -\frac{2e_a^2}{3} a_a^2 + e_a (A_{ba}^{\text{ret}} \cdot a_a) - \kappa_a (Q_{ab} \cdot a_a). \quad (5.17)$$

Similarly, the scalar product of particle's 4-velocity on the first order time derivative of particle's 4-momentum (5.14) is given by

$$(\dot{p}_a \cdot u_a) = \frac{2e_a^2}{3} \dot{a}_a^2 - (G_{ba}^{\text{ret}} \cdot u_a) - \kappa_a (\dot{Q}_{ab} \cdot u_a). \quad (5.18)$$

Equipped with the expression (5.7) one can derive the significant relation,

$$G_{ba}^\alpha = -F_{ba}^\alpha - e_a \frac{dA_{ba}^\alpha}{d\tau_a}, \quad (5.19)$$

where F_{ba}^α is the well-known Lorentz force of b -th charge acted on a -th one. Substituting this into eq.(5.18) and summing up (5.17) and modified (5.18), we obtain

$$(p_a \cdot u_a)^\cdot = e_a (A_{ba}^{\text{ret}} \cdot u_a)^\cdot - \kappa_a (Q_{ab} \cdot u_a)^\cdot \quad (5.20)$$

where dot means the a -th proper time derivative. On the other hand the scalar product of 4-momentum (5.16) on a -th 4-velocity is written as

$$(p_a \cdot u_a) = -m_a + e_a (A_{ba}^{\text{ret}} \cdot u_a) - \kappa_a (Q_{ab} \cdot u_a). \quad (5.21)$$

We see clearly that m_a is of constant value. It can be interpreted as already renormalized mass of a -th charged particle.

Finally, we differentiate the expression (5.16) and substitute it for the left-hand side of eq.(5.14). After cancellation of like terms and taking into account eq.(5.19) we arrive at the Lorentz-Dirac equation,

$$m_a a_a = \frac{2e_a^2}{3} (\dot{a}_a - a_a^2 u_a) + F_{ba}^{\text{ret}}. \quad (5.22)$$

We see that particles' equations of motion do not depend on mixing parameter κ .

In terms of kinematical variables conserved quantities of our particles plus field system looks as follows:

$$P = \sum_{a=1}^2 \left(m_a u_a - \frac{2e_a^2}{3} a_a + \frac{2e_a^2}{3} \int_{-\infty}^{\tau_a} ds_a a_a^2 u_a \right) - \sum_{a \neq b} \int_{-\infty}^{\tau_a} ds_a F_{ba}^{\text{ret}}, \quad (5.23)$$

$$\begin{aligned} \hat{M} &= \sum_{a=1}^2 \left[z_a \wedge \left(m_a u_a - \frac{2e_a^2}{3} a_a \right) + \frac{2e_a^2}{3} \int_{-\infty}^{\tau_a} ds_a (a_a^2 z_a \wedge u_a + u_a \wedge a_a) \right] \\ &\quad - \sum_{a \neq b} \int_{-\infty}^{\tau_a} ds_a z_a \wedge F_{ba}^{\text{ret}}. \end{aligned} \quad (5.24)$$

The work done by Lorentz forces of charges acting on one another exhausts the radiation reaction due to combination of fields.

Individual 4-momentum (5.16) of a -th dressed charged particle is modified comparing with the well-known Teitelboim's expression (1.7). The bound component $\hat{T}_{\text{int,bnd}}$ of mixed part of the electromagnetic field stress-energy tensor contributes two additional terms: “immovable core”, $e_a A_{ba}^{\text{ret}}$, of clear physical sense and “changeable shell”, $-\kappa_a Q_{ab}$, that heavily depends on mixing parameter κ . The corrections are inspired by unavoidable deformation of bound electromagnetic “clouds” due to mutual interaction between the sources. In my opinion, the changeable term arises due to forced choice of non-covariant surface of integration (see Subsection 4.4). For this reason we proclaim the expression

$$p_a = m_a u_a - \frac{2e_a^2}{3} a_a + e_a A_{ba}^{\text{ret}} \quad (5.25)$$

as the only one of true physical meaning.

6 Discrete symmetries

In the past Sections we have emphasized the importance of the invariance of the action integral (1.1) under the continuous group of space-time translations and rotations. According to Noether theorem, these symmetry properties imply conservation laws, i. e., those quantities that do not change with time. In this Section we study symmetry properties of energy-momentum and angular momentum carried by electromagnetic field which rely on invariance of (1.1) under discrete transformation groups.

6.1 Time inversion

The transformation of the time inversion $\mathcal{T}$ is defined by [2]

$$\mathcal{T} : y^\mu \mapsto y'^\mu = y_\mu. \quad (6.1)$$

It immediately gives

$$\mathcal{T} : z_a^\mu(s_a) \mapsto z_a'^\mu(s'_a) = z_{a,\mu}(s_a). \quad (6.2)$$

The proper time possesses odd parity [2],

$$\mathcal{T} : s_a \mapsto s'_a = -s_a. \quad (6.3)$$

The other kinematic quantities then follow easily,

$$\begin{aligned} \mathcal{T} : u_a^\mu(s_a) &\mapsto u_a'^\mu(s'_a) = -u_{a,\mu}(s_a), \\ \mathcal{T} : a_a^\mu(s_a) &\mapsto a_a'^\mu(s'_a) = a_{a,\mu}(s_a), \end{aligned} \quad (6.4)$$

etc. The retarded and the advanced instants transform into each other,

$$\mathcal{T} s_a^{\text{ret}}(s_b) = s_a^{\text{adv}}(s_b), \quad \mathcal{T} s_a^{\text{adv}}(s_b) = s_a^{\text{ret}}(s_b). \quad (6.5)$$

Inserting eqs. (6.4) and (6.5) into two-point function (4.36) yields

$$\begin{aligned}\mathcal{T}G_{ab}^\alpha[s_a^{ret}(s_b), s_b] &= -G_{ab,\alpha}[s_a^{adv}(s_b), s_b], \\ \mathcal{T}G_{ab}^\alpha[s_a^{adv}(s_b), s_b] &= -G_{ab,\alpha}[s_a^{ret}(s_b), s_b].\end{aligned}\tag{6.6}$$

To establish the symmetry properties of P_{rad} and $\hat{M}_{\text{rad}}$ with respect to time inversion, we locate the observation hyperplane at the distant future,

$$P_{\text{rad}} = \frac{2e_1^2}{3} \int_{-\infty}^{+\infty} ds_1 a_1^2 u_1 + \frac{2e_2^2}{3} \int_{-\infty}^{+\infty} ds_2 a_2^2 u_2 \tag{6.7}$$

$$\begin{aligned}&+ \kappa \int_{-\infty}^{+\infty} ds_1 (G_{21}^{\text{ret}} + G_{21}^{\text{adv}}) + (1 - \kappa) \int_{-\infty}^{+\infty} ds_2 (G_{12}^{\text{ret}} + G_{12}^{\text{adv}}), \\ \hat{M}_{\text{rad}} &= \frac{2e_1^2}{3} \int_{-\infty}^{+\infty} ds_1 (a_1^2 z_1 \wedge u_1 + u_1 \wedge a_1) + \frac{2e_2^2}{3} \int_{-\infty}^{+\infty} ds_2 (a_2^2 z_2 \wedge u_2 + u_2 \wedge a_2) \\ &+ \kappa \int_{-\infty}^{+\infty} ds_1 [z_1 \wedge (G_{21}^{\text{ret}} + G_{21}^{\text{adv}}) - e_1 u_1 \wedge (A_{21}^{\text{ret}} + A_{21}^{\text{adv}})] \\ &+ (1 - \kappa) \int_{-\infty}^{+\infty} ds_2 [z_2 \wedge (G_{12}^{\text{ret}} + G_{12}^{\text{adv}}) - e_2 u_2 \wedge (A_{12}^{\text{ret}} + A_{12}^{\text{adv}})].\end{aligned}\tag{6.8}$$

These expressions give the full amount of radiation emitted by interacting particles.

The transformation of the retarded and the advanced functions G_{ab} into each other implies that the radiated energy-momentum and angular momentum are of odd time parity,

$$\mathcal{T}P_{\text{rad}}^\mu = -P_{\text{rad},\mu}, \quad \mathcal{T}\hat{M}_{\text{rad}}^{\mu\nu} = -\hat{M}_{\text{rad},\mu\nu}.\tag{6.9}$$

6.2 Space inversion

This operation is defined by [2]

$$\mathcal{P} : y^\mu \mapsto y'^\mu = -y_\mu.\tag{6.10}$$

Correspondingly,

$$\mathcal{P} : z_a^\mu(s_a) \mapsto z_a'^\mu(s'_a) = -z_{a,\mu}(s_a).\tag{6.11}$$

The proper time s_a remains invariant. These transformation properties imply that that particles' 4-velocities and 4-accelerations change as follows:

$$\begin{aligned}\mathcal{P} : u_a^\mu(s_a) &\mapsto u_a'^\mu(s'_a) = -u_{a,\mu}(s_a), \\ \mathcal{P} : a_a^\mu(s_a) &\mapsto a_a'^\mu(s'_a) = -a_{a,\mu}(s_a).\end{aligned}\tag{6.12}$$

Since the retarded and the advanced instants remain invariant, the basic two-point function (4.36) transforms analogously,

$$\mathcal{P}G_{ab}^\alpha = -G_{ab,\alpha}.\tag{6.13}$$

Substituting this into eqs.(6.7) and (6.9) and using the relations (6.11) and (6.12) yields

$$\mathcal{P}P_{\text{rad}}^\mu = -P_{\text{rad},\mu}, \quad \mathcal{P}\hat{M}_{\text{rad}}^{\mu\nu} = \hat{M}_{\text{rad},\mu\nu}.\tag{6.14}$$

6.3 Reciprocity of particles 1 and 2

When a closed system of two identical charges is considered, radiative conserved quantities should be symmetric in indices 1 and 2 that label the particles. Having interchanged these indices in function (4.36) we obtain

$$\begin{aligned} G_{21}^\alpha[s_1, s_2^{ret}(s_1)]|_{1 \leftrightarrow 2} &= G_{12}^\alpha[s_1^{ret}(s_2), s_2], \\ G_{21}^\alpha[s_1, s_2^{adv}(s_1)]|_{1 \leftrightarrow 2} &= G_{12}^\alpha[s_1^{adv}(s_2), s_2]. \end{aligned} \quad (6.15)$$

From the reciprocity relations we see clearly that mixed parameter κ in expressions (6.7) and (6.9) should be equal to 1/2. Choosing the linear superposition

$$p_{\text{int,rad}} = \frac{1}{2} (p_{\text{rad},21} + p_{\text{rad},12}), \quad (6.16)$$

$$\hat{M}_{\text{int,rad}} = \frac{1}{2} (\hat{M}_{\text{rad},21} + \hat{M}_{\text{rad},12}), \quad (6.17)$$

we restore invariance of radiated energy-momentum and angular momentum with respect to reciprocity of particles 1 and 2.

7 Conclusions

The present paper is devoted to study of phenomena of emission and propagation of energy in classical electrodynamics. The field in action (1.1) has its own uncountably infinite degrees of freedom. Variation of (1.1) yields Maxwell's equations with point-like sources and equations of motions of particles interacting through the medium of the field. The problem then becomes one of mutual determination: the field is determined by the charged particles and their motion, and the motion of the charges is determined by the field.

In this paper we study interference of outgoing electromagnetic waves in a hyperplane $\Sigma_t = \{y \in \mathbb{M}_4 : y^0 = t\}$ associated with an unmoving inertial observer. We calculate how much electromagnetic field energy, momentum, and angular momentum flow across this hyperplane. Surface integration of the stress-energy tensor (1.4) over Σ_t reduces field's uncountably infinite degrees of freedom. After the renormalization procedure we arrive at the action at a distance theory [22, 23] where particles interact directly with one another. The fields in resulting expressions (5.23) and (5.24) do not have degrees of freedom of their own: they are functionals of particle paths. Following Ref. [24], we refer to them as direct particle fields.

Starting with the retarded Liénard-Wiechert fields, after integration we arrive at the retarded and the advanced direct particle fields. The retarded and the advanced instants arise naturally as the end points of interference integrals (2.15) and (2.16). But the retarded causality is not violated because the advanced instants as well as the retarded ones are before the fixed observation moment t . Direct particle fields referred to advanced instants do not describe neither incoming radiation nor converging electromagnetic waves.

Nevertheless, the retarded and the advanced quantities transform into each other under the influence of inversion of time axes. To demonstrate the invariance of electromagnetic

field's energy-momentum and angular momentum with respect to time inversion, we locate the observation surface Σ_t at distant future. We show that the full amount of radiation emitted by a closed system of two interacting charged particles is invariant with respect to past and future as well as with respect to inversion of space axes.

The part of Noether quantities which escapes to infinity is defined by basic two-point function (4.36). It is equal to the sum of Lorentz force F_{ab} of a -th charge acted on b -th one and the total time derivative of direct Liénard-Wiechert potential A_{ab} , taken with opposite sign (see eq.(5.19)). In the specific case of very massive particles, such that $e_1/m_1 \ll 1$ and $e_2/m_2 \ll 1$, the mixed energy-momentum $p_{\text{int,rad}}$ vanishes as could be expected for nonaccelerated charges. Indeed, let us consider static charge b at a coordinate origin. Zeroth component F_{ba}^0 , either retarded or advanced, is potential one: its integral over indicated portion of ζ_a cancels the change of only nontrivial A_{ba}^0 . Space components F_{ba}^i of the retarded and the advanced Lorentz forces compensate each other. Having performed a trivial Lorentz transformation we extend the statement on a motion with constant velocity. Radiative angular momentum possesses analogous properties.

To derive the equations of motion of interacting particles we compare flows of energy-momentum and angular momentum through very close hyperplanes Σ_t and $\Sigma_{t+\Delta t}$. Having balanced particles' individual characteristics and corresponding quantities carried by electromagnetic field, we obtain the well-known Lorentz-Dirac equation of motion of charged particle in the retarded field of the other charge where the self-action is taken into account. Since the Lorentz-Dirac equation possesses pathological solutions, such as runaway solutions or preaccelerations, the authors [25, 26, 27] propose the Landau-Lifshitz equation [9, §76] as more satisfactory alternative. Spohn [25] showed that a solution of the Lorentz-Dirac equation which does not satisfy the Landau-Lifshitz equation is of the runaway type. Moreover, Landau-Lifshitz equation does not permit runaway solutions or preacceleration [27].

Obtained expressions for radiated energy-momentum and angular momentum (including interference effects) are valuable also for a strictly prescribed motion of particles under the influence of a very powerful external force. It is necessary to compare them with corresponding results for circling charges [12, 13]. It would be interesting to consider the specific case when the charged particles are a very close to each other. Since the electromagnetic field satisfies the superposition principle, the models either an extended object consisting of N point charges or a continuous charge distribution are based on dynamics of two-body system [28, 29].

Acknowledgments

I am grateful to V.I. Tretyak for continuous encouragement and for a helpful reading of this manuscript. I would like to thank A. Duviryak for many useful discussions.

Appendix A Integration over angular variable

To calculate the mixed rates of energy-momentum (2.1) and angular momentum (2.2) carried by the electromagnetic field, we should first perform the integration over angle.

When facing this problem it is convenient to mark out φ -dependent terms in expressions under the integral sign. In the Maxwell energy-momentum tensor density we distinguish the second-order differential operator (3.10) with φ -dependent coefficients (3.7). It can be decomposed into a combination of partial derivatives in time variables $\hat{\Pi}_a$ given by eq.(3.13) and tail π_a of the type in eq.(3.14).

This Appendix is concerned with the computation of the tails. Equipped with them we express the aforementioned integrand as a combination of partial derivatives in t_1 and t_2 .

To implement this strategy we must first integrate the coefficients (3.7) over the angle variable. We start with the simplest one,

$$\mathcal{D}^a = \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{a}{qr_1 r_2}, \quad (\text{A.1})$$

where numerator a is equal to 1 or K_1^μ . Our task is to rewrite the integrand as a sum of term with denominator r_1 and term with denominator r_2 . To do it we introduce a new layer of mathematical formalism and develop convenient technique.

Let V be the vector space such that $\mathbf{i}_0, \mathbf{i}_1$, and $\mathbf{i}_2$ is its linear basis. We shall use $\eta_{\alpha\beta} = \text{diag}(-1, 1, 1)$ and its inverse $\eta^{\alpha\beta} = \text{diag}(-1, 1, 1)$ to lower and raise indices, respectively. We introduce the pairing

$$\begin{aligned} (\cdot) &: V \times V \rightarrow \mathbb{R} \\ (\mathbf{a} \cdot \mathbf{b}) &\mapsto \eta_{\alpha\beta} \mathbf{a}^\alpha \mathbf{b}^\beta, \end{aligned} \quad (\text{A.2})$$

which will be called the “scalar product”.

We introduce null vector $n^\alpha = (1, \sin \varphi, \cos \varphi)$ which belongs to the vector space V . We express the φ -dependent constructions

$$r_a = -(K_a \cdot v_a), \quad c_a = \gamma_a^{-2} + (K_a \cdot \dot{v}_a) \quad (\text{A.3})$$

as the scalar products $-(\mathbf{r}_a \cdot n)$ and $(\mathbf{c}_a \cdot n)$, respectively. We shall use **sans-serif** letters for the components of timelike three-vectors $\mathbf{r}_a \in V$ and $\mathbf{c}_a \in V$,

$$\mathbf{r}_a^0 = k_a^0 - (\mathbf{v}_a \mathbf{n}_q) k_a^3, \quad \mathbf{r}_a^1 = h v_a^i \omega_i^1, \quad \mathbf{r}_a^2 = h v_a^i \omega_i^2; \quad (\text{A.4})$$

$$\mathbf{c}_a^0 = -\gamma_a^{-2} - (\dot{\mathbf{v}}_a \mathbf{n}_q) k_a^3, \quad \mathbf{c}_a^1 = h \dot{v}_a^i \omega_i^1, \quad \mathbf{c}_a^2 = h \dot{v}_a^i \omega_i^2, \quad (\text{A.5})$$

where orthogonal matrix,

$$\hat{\omega} = \begin{pmatrix} \cos \varphi_q & -\sin \varphi_q & 0 \\ \sin \varphi_q & \cos \varphi_q & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \cos \vartheta_q & 0 & \sin \vartheta_q \\ 0 & 1 & 0 \\ -\sin \vartheta_q & 0 & \cos \vartheta_q \end{pmatrix}, \quad (\text{A.6})$$

is constructed from components of the relative position 3-vector $\mathbf{q} = \mathbf{z}_1 - \mathbf{z}_2$. The numerator in eq.(A.1) is the scalar product $(\mathbf{a} \cdot n) = -\mathbf{a}^0 + \mathbf{a}^1 \sin \varphi + \mathbf{a}^2 \cos \varphi$; it is equal to 1 if vector $\mathbf{a} = (-1, 0, 0)$.

We introduce the dual space of one-forms, say W , with basis $\hat{\omega}^0, \hat{\omega}^1$ and $\hat{\omega}^2$, such that $\hat{\omega}^\mu(\mathbf{i}_\nu) = \delta_\nu^\mu$, where $\mathbf{i}_0, \mathbf{i}_1, \mathbf{i}_2$ constitute the basis of V . The wedge product $\hat{L} = \hat{a} \wedge \hat{b}$ of two one forms $\hat{a}$ and $\hat{b}$ constitutes two-form,

$$\hat{L} = (a_0 b_1 - a_1 b_0) \hat{\omega}^0 \wedge \hat{\omega}^1 + (a_0 b_2 - a_2 b_0) \hat{\omega}^0 \wedge \hat{\omega}^2 + (a_1 b_2 - a_2 b_1) \hat{\omega}^1 \wedge \hat{\omega}^2. \quad (\text{A.7})$$

We introduce dual three-vector $\mathbf{L} = {}^* \hat{L}$ with components

$$\begin{aligned} \mathbf{L}^\alpha &= \frac{1}{2!} \varepsilon^{\alpha\beta\gamma} L_{\beta\gamma} \\ &= \varepsilon^{\alpha\beta\gamma} a_\beta b_\gamma. \end{aligned} \quad (\text{A.8})$$

$\varepsilon^{\alpha\beta\gamma}$ denotes the Ricci symbol in three dimensions,

$$\varepsilon^{\alpha\beta\gamma} = \begin{cases} 1 & \text{when } \alpha\beta\gamma \text{ is an even permutation of } 0, 1, 2 \\ -1 & \text{when } \alpha\beta\gamma \text{ is an odd permutation of } 0, 1, 2 \\ 0 & \text{otherwise.} \end{cases} \quad (\text{A.9})$$

We raise indices in eq.(A.8) and define the vector product of two vectors, $\mathbf{a}$ and $\mathbf{b}$,

$$\mathbf{L}^\alpha = \varepsilon^\alpha_{\mu\nu} \mathbf{a}^\mu \mathbf{b}^\nu. \quad (\text{A.10})$$

Tensor,

$$\varepsilon^\alpha_{\mu\nu} = \varepsilon^{\alpha\beta\gamma} \eta_{\beta\mu} \eta_{\gamma\nu}, \quad (\text{A.11})$$

has the components

$$\varepsilon^0_{\mu\nu} = \varepsilon^{0\mu\nu}, \quad \varepsilon^1_{\mu\nu} = -\varepsilon^{1\mu\nu}, \quad \varepsilon^2_{\mu\nu} = -\varepsilon^{2\mu\nu}. \quad (\text{A.12})$$

Now we calculate the double vector product,

$$[\mathbf{A}[\mathbf{BC}]]^\alpha = \varepsilon^\alpha_{\beta\gamma} \mathbf{A}^\beta \varepsilon^\gamma_{\mu\nu} \mathbf{B}^\mu \mathbf{C}^\nu. \quad (\text{A.13})$$

Since

$$\varepsilon^\alpha_{\beta\gamma} \varepsilon^\gamma_{\mu\nu} = -\delta^\alpha_\mu \eta_{\beta\nu} + \delta^\alpha_\nu \eta_{\beta\mu}, \quad (\text{A.14})$$

we arrive to the unusual rule,

$$[\mathbf{A}[\mathbf{BC}]] = -\mathbf{B}(\mathbf{A} \cdot \mathbf{C}) + \mathbf{C}(\mathbf{A} \cdot \mathbf{B}), \quad (\text{A.15})$$

instead of the well-known law acting in space with Euclidean metric.

To simplify the denominator $r_1 r_2$ in the integrand of eq.(A.1) as much as possible, we rewrite 2π -periodic functions $r_a = -r_{a,0} - r_{a,1} \sin \varphi - r_{a,2} \cos \varphi$ as follows:

$$r_a = -r_{a,0} - \rho_a \sin(\varphi + \phi_a), \quad \rho_a = \sqrt{r_{a,1}^2 + r_{a,2}^2}. \quad (\text{A.16})$$

(We recall that r_a is the scalar products $(\mathbf{r}_a \cdot \mathbf{n})$ taken with opposite sign, components r_a^μ are given by eqs.(A.4).) Shift in argument of harmonic function is determined by the relations

$$r_{a,1} = \rho_a \cos \phi_a, \quad r_{a,2} = \rho_a \sin \phi_a. \quad (\text{A.17})$$

After some algebra we rewrite the integrand of eq.(A.1) as the following sum:

$$\frac{a}{r_1 r_2} = \frac{A_{12}^a + C_{12}^a \rho_1 \cos(\varphi + \phi_1)}{r_1} + \frac{A_{21}^a - C_{12}^a \rho_2 \cos(\varphi + \phi_2)}{r_2}, \quad (\text{A.18})$$

where $a = (\mathbf{a} \cdot \mathbf{n})$. Coefficients A_{12}^a, A_{21}^a and C_{12}^a satisfy the vector equation

$$-A_{12}^a \mathbf{r}_2 - A_{21}^a \mathbf{r}_1 + C_{12}^a \mathbf{L}_{12} = \mathbf{a}, \quad (\text{A.19})$$

where by $\mathbf{r}_1$ and $\mathbf{r}_2$ we mean three-vectors with components in eq.(A.4) and $\mathbf{L}_{12} = [\mathbf{r}_1 \mathbf{r}_2]$.

To solve equation (A.19) we postmultiply it on the vector product $[\mathbf{r}_1 \mathbf{L}_{12}]$, then on the vector product $[\mathbf{r}_2 \mathbf{L}_{21}]$, and, finally, on $\mathbf{L}_{12}$. After some algebra we obtain

$$A_{12}^a = \frac{([\mathbf{a} \mathbf{r}_1] \cdot \mathbf{L}_{12})}{D_{12}}, \quad A_{21}^a = \frac{([\mathbf{a} \mathbf{r}_2] \cdot \mathbf{L}_{21})}{D_{21}}, \quad C_{12}^a = \frac{(\mathbf{a} \cdot \mathbf{L}_{12})}{D_{12}}, \quad (\text{A.20})$$

where the denominator $D_{12} = (\mathbf{L}_{12} \cdot \mathbf{L}_{12})$ is symmetric in its indices.

Substituting eq.(A.18) into eq.(A.1) and using the identities

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} \frac{d\varphi}{r_a^0 - \rho_a \sin(\varphi + \phi_a)} &= \frac{1}{\sqrt{(r_a^0)^2 - \rho_a^2}} \\ &= \frac{1}{\|\mathbf{r}_a\|}, \\ \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{\cos(\varphi + \phi_a)}{r_a^0 - \rho_a \sin(\varphi + \phi_a)} &= 0 \end{aligned} \quad (\text{A.21})$$

yields

$$\mathcal{D}^a = \frac{A_{12}^a}{q\|\mathbf{r}_1\|} + \frac{A_{21}^a}{q\|\mathbf{r}_2\|} \quad (\text{A.22})$$

after integration over φ .

Now we turn to the calculation of the coefficient

$$\mathcal{B}^a = \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{ac_2}{qr_1(r_2)^2}. \quad (\text{A.23})$$

Equipped with the relations in eq.(A.18) we rewrite the integrand as a sum of terms which are proportional to the $1/r_1$, $1/r_2$, and $1/(r_2)^2$, respectively. Using the identities

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} \frac{d\varphi}{[r_a^0 - \rho_a \sin(\varphi + \phi_a)]^2} &= \frac{r_a^0}{\|\mathbf{r}_a\|^3}, \\ \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{\cos(\varphi + \phi_a)}{[r_a^0 - \rho_a \sin(\varphi + \phi_a)]^2} &= 0 \end{aligned} \quad (\text{A.24})$$

and taking into account the relations in eq.(A.21) gives

$$\begin{aligned} \mathcal{B}^a &= -\frac{1}{q\|\mathbf{r}_2\|^3} \frac{(\mathbf{a} \cdot \mathbf{r}_2)(\mathbf{c}_2 \cdot \mathbf{r}_2)}{D_{21}} (\mathbf{r}_2 \cdot \mathbf{r}_1) + \frac{1}{q\|\mathbf{r}_2\|} \left[A_{12}^{c_2} A_{21}^a + A_{12}^a A_{21}^{c_2} - \frac{(\mathbf{a} \cdot \mathbf{c}_2)(\mathbf{r}_1 \cdot \mathbf{r}_2)}{D_{21}} \right] \\ &+ \frac{1}{q\|\mathbf{r}_1\|} \left[2A_{12}^a A_{12}^{c_2} - \frac{([\mathbf{a} \mathbf{r}_1] \cdot [\mathbf{c}_2 \mathbf{r}_1])}{D_{12}} \right]. \end{aligned} \quad (\text{A.25})$$

The resulting expression for the term

$$\mathcal{C}^a = \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{ac_1}{q(r_1)^2 r_2} \quad (\text{A.26})$$

can be obtained by interchanging indices 1 and 2 in the right-hand side of eq.(A.25).

After a routine computation based on the repeated usage of relation (A.18), we find the most complicate term,

$$\begin{aligned} \mathcal{A}^a &= \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{ac_1 c_2}{q(r_1)^2 (r_2)^2} \\ &= \frac{B_{12}}{q\|r_1\|} + \frac{B_{21}}{q\|r_2\|} + J_1 \frac{(\mathbf{a} \cdot \mathbf{r}_1)}{q\|r_1\|^3} + J_2 \frac{(\mathbf{a} \cdot \mathbf{r}_2)}{q\|r_2\|^3}, \end{aligned} \quad (\text{A.27})$$

where

$$\begin{aligned} J_1 &= 2A_{12}^{c_1} A_{12}^{c_2} - \frac{([\mathbf{c}_1 \mathbf{r}_1] \cdot [\mathbf{c}_2 \mathbf{r}_1])}{D_{12}}, \\ B_{12} &= 3A_{12}^a A_{12}^{c_1} A_{21}^{c_2} + 3A_{12}^a A_{12}^{c_2} A_{21}^{c_1} + 2A_{12}^{c_1} A_{12}^{c_2} A_{21}^a + A_{12}^a \left\{ \frac{([\mathbf{c}_1 \mathbf{r}_1] \cdot [\mathbf{c}_2 \mathbf{r}_2])}{D_{12}} + \frac{([\mathbf{c}_1 \mathbf{r}_2] \cdot [\mathbf{c}_2 \mathbf{r}_1])}{D_{12}} \right\} \\ &\quad - A_{21}^{c_1} \frac{([\mathbf{c}_2 \mathbf{r}_1] \cdot [\mathbf{a} \mathbf{r}_1])}{D_{12}} - A_{21}^{c_2} \frac{([\mathbf{c}_1 \mathbf{r}_1] \cdot [\mathbf{a} \mathbf{r}_1])}{D_{12}} + A_{12}^{c_1} \frac{([\mathbf{c}_2 \mathbf{r}_1] \cdot [\mathbf{a} \mathbf{r}_2])}{D_{12}} + A_{12}^{c_2} \frac{([\mathbf{c}_1 \mathbf{r}_1] \cdot [\mathbf{a} \mathbf{r}_2])}{D_{12}}, \end{aligned} \quad (\text{A.28})$$

and the others, B_{21} and J_2 , can be obtained via interchanging indices 1 and 2.

We now turn to the differentiation of coefficient (A.22) with respect to time variables t_1 and t_2 . Having substituted $\mathbf{o} = (-1, 0, 0)$ for $\mathbf{a}$ in the expressions (A.22), (A.25), and (A.27) we obtain the terms $\mathcal{D}^0$, $\mathcal{B}^0$ and $\mathcal{A}^0$, respectively. The remaining term, $\mathcal{C}^0$, can be obtained from $\mathcal{B}^0$ via reciprocity. The calculations are based on the relations obtained via differentiation of third components k_a^3 and h^2 , i.e. the square of radius of the circle $C(O, h) = S_1 \cap S_2$,

$$\begin{aligned} \frac{\partial k_a^3}{\partial t_a} &= - \left[(-1)^a \frac{r_a^0}{q} + (\mathbf{n}_q \mathbf{v}_a) \right], \quad \frac{\partial k_a^3}{\partial t_b} = (-1)^a \frac{r_b^0}{q}, \\ \frac{\partial h^2}{\partial t_1} &= -2k_2^3 \frac{r_1^0}{q}, \quad \frac{\partial h^2}{\partial t_2} = 2k_1^3 \frac{r_2^0}{q}. \end{aligned} \quad (\text{A.29})$$

They immediately give

$$\begin{aligned} \frac{\partial r_a^0}{\partial t_a} &= c_a^0 + \frac{(-1)^a}{q} \left[(\mathbf{v}_a \mathbf{n}_q) r_a^0 + [\mathbf{v}_a \mathbf{n}_q]^2 k_b^3 \right], \\ \frac{\partial r_a^0}{\partial t_b} &= \frac{(-1)^b}{q} \left[(\mathbf{v}_a \mathbf{n}_q) r_b^0 + ([\mathbf{v}_a \mathbf{n}_q][\mathbf{v}_b \mathbf{n}_q]) k_a^3 \right], \end{aligned} \quad (\text{A.30})$$

and eventually give

$$\begin{aligned} \frac{\partial(\mathbf{r}_a \cdot \mathbf{r}_a)}{\partial t_a} &= 2 \left[(\mathbf{r}_a \cdot \mathbf{c}_a) + (-1)^a \frac{(\mathbf{v}_a \mathbf{n}_q)}{q} (\mathbf{r}_a \cdot \mathbf{r}_a) \right], \\ \frac{\partial(\mathbf{r}_a \cdot \mathbf{r}_a)}{\partial t_b} &= 2(-1)^b \left[\frac{(\mathbf{v}_a \mathbf{n}_q)}{q} (\mathbf{r}_a \cdot \mathbf{r}_b) + \frac{k_b^3 r_b^0 (\mathbf{r}_a \cdot \mathbf{r}_a) - r_a^0 (\mathbf{r}_a \cdot \mathbf{r}_b)}{h^2} \right], \\ \frac{\partial(\mathbf{r}_b \cdot \mathbf{r}_a)}{\partial t_b} &= (\mathbf{r}_a \cdot \mathbf{c}_b) + (-1)^b \left[\frac{(\mathbf{v}_b \mathbf{n}_q)}{q} (\mathbf{r}_b \cdot \mathbf{r}_a) + \frac{(\mathbf{v}_a \mathbf{n}_q)}{q} (\mathbf{r}_b \cdot \mathbf{r}_b) + \frac{k_a^3 r_b^0 (\mathbf{r}_b \cdot \mathbf{r}_a) - r_a^0 (\mathbf{r}_b \cdot \mathbf{r}_b)}{h^2} \right]. \end{aligned} \quad (\text{A.31})$$

We use Latin indices a and b which run from 1 to 2 ($a \neq b$). We use bold script for conventional three-velocities $v_a^i = dz_a^i/dt_a$ and unit three-vector $n_q^i = q^i/q$ in $\mathbf{q}$ -direction. By $(\mathbf{v}_b \mathbf{n}_q)$ and $[\mathbf{v}_b \mathbf{n}_q]$ we denote the conventional scalar product and cross product of these vectors, respectively.

Usage of these relations allows us to calculate the derivatives of coefficients A_{ab}^0 :

$$\begin{aligned} \frac{\partial A_{ab}^0}{\partial t_a} &= A_{ab}^0 A_{ba}^{c_a} + A_{ab}^{c_a} A_{ba}^0 + \frac{([\mathbf{or}_a] \cdot [\mathbf{c}_a \mathbf{r}_b])}{D_{ab}}, \\ \frac{\partial A_{ab}^0}{\partial t_b} &= 2A_{ab}^{c_b} A_{ab}^0 - \frac{([\mathbf{or}_a] \cdot [\mathbf{c}_b \mathbf{r}_a])}{D_{ab}} + (-1)^b \frac{k_a^3}{qh^2} (\mathbf{o}^0 + \mathbf{r}_2^0 A_{12}^0 + \mathbf{r}_1^0 A_{21}^0) \\ &\quad + (-1)^a \left(A_{ab}^0 \frac{(\mathbf{n}_q \mathbf{v}_b)}{q} + A_{ba}^0 \frac{(\mathbf{n}_q \mathbf{v}_a)}{q} \right). \end{aligned} \quad (\text{A.32})$$

Substituting these into equality

$$\frac{\partial \mathcal{D}^0}{\partial t_a} = \frac{\partial}{\partial t_a} \left(\frac{A_{12}^0}{q\|\mathbf{r}_1\|} + \frac{A_{21}^0}{q\|\mathbf{r}_2\|} \right) \quad (\text{A.33})$$

and using the identities

$$\begin{aligned} \frac{\partial}{\partial t_a} \left(\frac{1}{\|\mathbf{r}_a\|} \right) &= \frac{(\mathbf{c}_a \cdot \mathbf{r}_a)}{\|\mathbf{r}_a\|^3} - (-1)^a \frac{1}{\|\mathbf{r}_a\|} \frac{(\mathbf{n}_q \mathbf{v}_a)}{q}, \\ \frac{\partial}{\partial t_b} \left(\frac{1}{\|\mathbf{r}_a\|} \right) &= (-1)^b \frac{(\mathbf{r}_a \cdot \mathbf{r}_b)}{\|\mathbf{r}_a\|^3} \frac{(\mathbf{n}_q \mathbf{v}_a)}{q} + (-1)^a \frac{k_a^3}{q\|\mathbf{r}_a\|^3} \frac{\mathbf{r}_a^0 (\mathbf{r}_a \cdot \mathbf{r}_b) - \mathbf{r}_b^0 (\mathbf{r}_a \cdot \mathbf{r}_a)}{h^2} \end{aligned} \quad (\text{A.34})$$

yields

$$\frac{\partial \mathcal{D}^0}{\partial t_1} = \mathcal{C}^0 + \frac{k_2^0 (\mathbf{n}_q \mathbf{v}_2) - k_2^3 \mathbf{v}_2^2}{q^2 \|\mathbf{r}_2\|^3}, \quad \frac{\partial \mathcal{D}^0}{\partial t_2} = \mathcal{B}^0 - \frac{k_1^0 (\mathbf{n}_q \mathbf{v}_1) - k_1^3 \mathbf{v}_1^2}{q^2 \|\mathbf{r}_1\|^3}. \quad (\text{A.35})$$

Further we calculate the partial derivative $\partial \mathcal{C}^0 / \partial t_2$, subtract it from $\mathcal{A}^0$, and prove the identity

$$\mathcal{A}^0 - \frac{\partial \mathcal{C}^0}{\partial t_2} = \frac{\partial}{\partial t_1} \left(\mathcal{B}^0 - \frac{\partial \mathcal{D}^0}{\partial t_2} \right) \quad \text{i.e.} \quad \pi^0 = 0. \quad (\text{A.36})$$

(One can derive $\partial \mathcal{B}^0 / \partial t_1$, subtract it from $\mathcal{A}^0$, and compare the result with $\partial / \partial t_2 (\mathcal{C}^0 - \partial \mathcal{D}^0 / \partial t_1)$.)

Now, we calculate the tail

$$\pi_a^\alpha = \mathcal{A}_a^\alpha - \frac{\partial \mathcal{B}_a^\alpha}{\partial t_1} - \frac{\partial \mathcal{C}_a^\alpha}{\partial t_2} + \frac{\partial^2 \mathcal{D}_a^\alpha}{\partial t_1 \partial t_2}, \quad (\text{A.37})$$

where

$$\begin{aligned} \mathcal{D}_a^\alpha &= \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{K_a^\alpha}{q r_1 r_2}, & \mathcal{B}_a^\alpha &= \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{K_a^\alpha c_2}{q r_1 (r_2)^2}, \\ \mathcal{C}_a^\alpha &= \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{K_a^\alpha c_1}{q (r_1)^2 r_2}, & \mathcal{A}_a^\alpha &= \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{K_a^\alpha c_1 c_2}{q (r_1)^2 (r_2)^2}. \end{aligned} \quad (\text{A.38})$$

The zeroth component, $K_a^0 = k_a^0$, does not depend on φ . Inserting relations $\mathcal{D}_a^0 = k_a^0 \mathcal{D}^0$, $\mathcal{B}_a^0 = k_a^0 \mathcal{B}^0$, $\mathcal{C}_a^0 = k_a^0 \mathcal{C}^0$, and $\mathcal{A}_a^0 = k_a^0 \mathcal{A}^0$ into eq.(A.37) and taking into account identity (A.36) yields

$$\pi_1^0 = \mathcal{B}^0 - \frac{\partial \mathcal{D}^0}{\partial t_2}, \quad \pi_2^0 = \mathcal{C}^0 - \frac{\partial \mathcal{D}^0}{\partial t_1}. \quad (\text{A.39})$$

Space components, K_a^i , depend on φ . They can be expressed as the scalar product $(\mathbf{K}_a^i \cdot \mathbf{n})$ where components of three-vectors $\mathbf{K}_a^i \in V$ are as follows:

$$\mathbf{K}_{a,0}^i = n_q^i k_a^3, \quad \mathbf{K}_{a,1}^i = h \omega^i{}_1, \quad \mathbf{K}_{a,2}^i = h \omega^i{}_2. \quad (\text{A.40})$$

Here $\omega^i{}_j$ are components of the orthogonal matrix (A.6). Having substituted $\mathbf{K}_a^i$ for $\mathbf{a}$ in expressions (A.22), (A.25), and (A.27) we obtain the terms $\mathcal{D}_a^i$, $\mathcal{B}_a^i$ and $\mathcal{A}_a^i$, respectively. The last term, $\mathcal{C}_a^i$, can be obtained from $\mathcal{B}_a^i$ via reciprocity. To differentiate them we need the equalities

$$\begin{aligned} \frac{\partial(\mathbf{K}_1^i \cdot \mathbf{r}_1)}{\partial t_1} &= (\mathbf{K}_1^i \cdot \mathbf{c}_1) - v_1^i r_1^0 - \frac{(\mathbf{n}_q \mathbf{v}_1)}{q} (\mathbf{K}_1^i \cdot \mathbf{r}_1) - \frac{n_q^i}{q} (\mathbf{r}_1 \cdot \mathbf{r}_1) \\ &\quad - \frac{k_2^3}{qh^2} [\mathbf{K}_{1,0}^i (\mathbf{r}_1 \cdot \mathbf{r}_1) + r_1^0 (\mathbf{K}_1^i \cdot \mathbf{r}_1)], \\ \frac{\partial(\mathbf{K}_1^i \cdot \mathbf{r}_2)}{\partial t_1} &= -v_1^i r_2^0 - \frac{(\mathbf{n}_q \mathbf{v}_2)}{q} (\mathbf{K}_1^i \cdot \mathbf{r}_1) - \frac{n_q^i}{q} (\mathbf{r}_1 \cdot \mathbf{r}_2) \\ &\quad + \frac{k_2^3}{qh^2} [r_2^0 (\mathbf{K}_1^i \cdot \mathbf{r}_1) - \mathbf{K}_{1,0}^i (\mathbf{r}_1 \cdot \mathbf{r}_2) - 2r_1^0 (\mathbf{K}_1^i \cdot \mathbf{r}_2)], \\ \frac{\partial(\mathbf{K}_1^i \cdot \mathbf{r}_1)}{\partial t_2} &= \frac{(\mathbf{n}_q \mathbf{v}_1)}{q} (\mathbf{K}_1^i \cdot \mathbf{r}_2) + \frac{n_q^i}{q} (\mathbf{r}_1 \cdot \mathbf{r}_2) \\ &\quad + \frac{k_1^3}{qh^2} [-r_1^0 (\mathbf{K}_1^i \cdot \mathbf{r}_2) + \mathbf{K}_{1,0}^i (\mathbf{r}_1 \cdot \mathbf{r}_2) + 2r_2^0 (\mathbf{K}_1^i \cdot \mathbf{r}_1)], \\ \frac{\partial(\mathbf{K}_1^i \cdot \mathbf{r}_2)}{\partial t_2} &= (\mathbf{K}_1^i \cdot \mathbf{c}_2) + \frac{(\mathbf{n}_q \mathbf{v}_2)}{q} (\mathbf{K}_1^i \cdot \mathbf{r}_2) + \frac{n_q^i}{q} (\mathbf{r}_2 \cdot \mathbf{r}_2) \\ &\quad + \frac{k_1^3}{qh^2} [\mathbf{K}_{1,0}^i (\mathbf{r}_2 \cdot \mathbf{r}_2) + r_2^0 (\mathbf{K}_1^i \cdot \mathbf{r}_2)], \end{aligned}$$

in addition to eqs.(A.31) and (A.34).

The derivation of equalities

$$\begin{aligned} \mathcal{C}_1^i - \frac{\partial \mathcal{D}_1^i}{\partial t_1} &= v_1^i \mathcal{D}^0 - \frac{n_q^i}{q^2 \|\mathbf{r}_2\|} - \frac{1}{q \|\mathbf{r}_2\|^3} \frac{(\mathbf{n}_q \mathbf{v}_2)}{q} (\mathbf{K}_1^i \cdot \mathbf{r}_2) \\ &\quad + \frac{1}{q \|\mathbf{r}_2\|^3} \frac{k_2^3}{q} \left(\mathbf{K}_{1,0}^i [\mathbf{n}_q \mathbf{v}_2]^2 + r_2^0 [\mathbf{n}_q [\mathbf{v}_2 \mathbf{n}_q]]^i \right), \\ \mathcal{B}_1^i - \frac{\partial \mathcal{D}_1^i}{\partial t_2} &= \frac{n_q^i}{q^2 \|\mathbf{r}_1\|} + \frac{1}{q \|\mathbf{r}_1\|^3} \frac{(\mathbf{n}_q \mathbf{v}_1)}{q} (\mathbf{K}_1^i \cdot \mathbf{r}_1) \\ &\quad - \frac{1}{q \|\mathbf{r}_1\|^3} \frac{k_1^3}{q} \left(\mathbf{K}_{1,0}^i [\mathbf{n}_q \mathbf{v}_1]^2 + r_1^0 [\mathbf{n}_q [\mathbf{v}_1 \mathbf{n}_q]]^i \right), \end{aligned} \quad (\text{A.41})$$

is virtually identical to that presented above, and we shall not bother with the details. By virtue of the relation $K_2^i - K_1^i = q^i$ the integrals (A.38) are related as follows:

$$\mathcal{D}_2^i = \mathcal{D}_1^i + q^i \mathcal{D}^0, \quad \mathcal{B}_2^i = \mathcal{B}_1^i + q^i \mathcal{B}^0, \quad \mathcal{C}_2^i = \mathcal{C}_1^i + q^i \mathcal{C}^0, \quad \mathcal{A}_2^i = \mathcal{A}_1^i + q^i \mathcal{A}^0. \quad (\text{A.42})$$

Equipped with these expressions we find

$$\begin{aligned} \mathcal{C}_2^i - \frac{\partial \mathcal{D}_2^i}{\partial t_1} &= -\frac{n_q^i}{q^2 \|r_2\|} - \frac{1}{q \|r_2\|^3} \frac{(\mathbf{n}_q \mathbf{v}_2)}{q} (\mathbf{K}_2^i \cdot \mathbf{r}_2) \\ &\quad + \frac{1}{q \|r_2\|^3} \frac{k_2^3}{q} \left(\mathbf{K}_{2,0}^i [\mathbf{n}_q \mathbf{v}_2]^2 + r_2^0 [\mathbf{n}_q [\mathbf{v}_2 \mathbf{n}_q]]^i \right), \\ \mathcal{B}_2^i - \frac{\partial \mathcal{D}_2^i}{\partial t_2} &= v_2^i \mathcal{D}^0 + \frac{n_q^i}{q^2 \|r_1\|} + \frac{1}{q \|r_1\|^3} \frac{(\mathbf{n}_q \mathbf{v}_1)}{q} (\mathbf{K}_2^i \cdot \mathbf{r}_1) \\ &\quad - \frac{1}{q \|r_1\|^3} \frac{k_1^3}{q} \left(\mathbf{K}_{2,0}^i [\mathbf{n}_q \mathbf{v}_1]^2 + r_1^0 [\mathbf{n}_q [\mathbf{v}_1 \mathbf{n}_q]]^i \right). \end{aligned} \quad (\text{A.43})$$

Finally, after a straightforward (but fairly lengthy) calculations we derive the following relations:

$$\pi_1^i = v_1^i \left(\mathcal{B}^0 - \frac{\partial \mathcal{D}^0}{\partial t_2} \right), \quad \pi_2^i = v_2^i \left(\mathcal{C}^0 - \frac{\partial \mathcal{D}^0}{\partial t_1} \right), \quad (\text{A.44})$$

which generalize eqs.(A.39).

We will need also the tail

$$\pi_{12}^{\alpha\beta} = \frac{\partial^2 \mathcal{D}_{12}^{\alpha\beta}}{\partial t_1 \partial t_2} - \frac{\partial \mathcal{B}_{12}^{\alpha\beta}}{\partial t_1} - \frac{\partial \mathcal{C}_{12}^{\alpha\beta}}{\partial t_2} + \mathcal{A}_{12}^{\alpha\beta}, \quad (\text{A.45})$$

where

$$\begin{aligned} \mathcal{D}_{12}^{\alpha\beta} &= \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{K_1^\alpha K_2^\beta}{q r_1 r_2}, & \mathcal{B}_{12}^{\alpha\beta} &= \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{K_1^\alpha K_2^\beta c_2}{q r_1 (r_2)^2}, \\ \mathcal{C}_{12}^{\alpha\beta} &= \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{K_1^\alpha K_2^\beta c_1}{q (r_1)^2 r_2}, & \mathcal{A}_{12}^{\alpha\beta} &= \frac{1}{2\pi} \int_0^{2\pi} d\varphi \frac{K_1^\alpha K_2^\beta c_1 c_2}{q (r_1)^2 (r_2)^2}. \end{aligned} \quad (\text{A.46})$$

It can be obtained by means of covariant generalization of previous relations. Setting $\alpha = 0$ and $\beta = 0$ and taking into account eq.(A.36), we obtain

$$\begin{aligned} \pi_{12}^{00} &= k_1^0 \left(\mathcal{C}^0 - \frac{\partial \mathcal{D}^0}{\partial t_1} \right) + k_2^0 \left(\mathcal{B}^0 - \frac{\partial \mathcal{D}^0}{\partial t_2} \right) + \mathcal{D}^0 \\ &= \mathcal{C}_1^0 - \frac{\partial \mathcal{D}_1^0}{\partial t_1} + \mathcal{B}_2^0 - \frac{\partial \mathcal{D}_2^0}{\partial t_2} - \mathcal{D}^0, \end{aligned} \quad (\text{A.47})$$

where relations $\mathcal{D}_a^0 = k_a^0 \mathcal{D}^0$, $\mathcal{B}_a^0 = k_a^0 \mathcal{B}^0$, and $\mathcal{C}_a^0 = k_a^0 \mathcal{C}^0$ are taken into account. If $\alpha = i$ and $\beta = 0$, we have

$$\begin{aligned} \pi_{12}^{i0} &= \mathcal{C}_1^i - \frac{\partial \mathcal{D}_1^i}{\partial t_1} + k_2^0 v_1^i \left(\mathcal{B}^0 - \frac{\partial \mathcal{D}^0}{\partial t_2} \right) \\ &= \mathcal{C}_1^i - \frac{\partial \mathcal{D}_1^i}{\partial t_1} + v_1^i \left(\mathcal{B}_2^0 - \frac{\partial \mathcal{D}_2^0}{\partial t_2} \right) - v_1^i \mathcal{D}^0. \end{aligned} \quad (\text{A.48})$$

If $\alpha = 0$ and $\beta = j$, we arrive at

$$\begin{aligned}\pi_{12}^{0j} &= k_1^0 v_2^j \left(\mathcal{C}^0 - \frac{\partial \mathcal{D}^0}{\partial t_1} \right) + \mathcal{B}_2^j - \frac{\partial \mathcal{D}_2^j}{\partial t_2} \\ &= v_2^j \left(\mathcal{C}_1^0 - \frac{\partial \mathcal{D}_1^0}{\partial t_1} \right) + \mathcal{B}_2^j - \frac{\partial \mathcal{D}_2^j}{\partial t_2} - v_2^j \mathcal{D}^0.\end{aligned}\quad (\text{A.49})$$

An obvious generalization of expressions (A.47)-(A.49) is

$$\pi_{12}^{\alpha\beta} = v_1^\alpha \left(\mathcal{B}_2^\beta - \frac{\partial \mathcal{D}_2^\beta}{\partial t_2} \right) + v_2^\beta \left(\mathcal{C}_1^\alpha - \frac{\partial \mathcal{D}_1^\alpha}{\partial t_1} \right) - v_1^\alpha v_2^\beta \mathcal{D}^0. \quad (\text{A.50})$$

Since the angular integration leads to a combination of partial derivatives in time variables, the end points are valuable only. At these points the radius of circle $C(O, h) = S_1 \cap S_2$ vanishes. Hence we can restrict ourselves to calculation of expressions $\mathcal{B}_{12}^{ij} - \partial_2 \mathcal{D}_{12}^{ij}$ and $\mathcal{C}_{12}^{ij} - \partial_1 \mathcal{D}_{12}^{ij}$ at points where $h = 0$.

To simplify the calculations as much as possible we express the integrands of eqs.(A.46) in form of expansions in powers of h . It allows us to remove harmonic functions from denominators. Since the derivatives $\partial h^2 / \partial t_a$ do not vanish whenever $h^2 = 0$ (see eq.(A.29)), it is sufficient to expand $\mathcal{D}_{12}^{ij}$ up to the first order of this parameter,

$$\begin{aligned}\mathcal{D}_{12}^{ij} &= \frac{k_1^3 k_2^3 n_q^i n_q^j}{q r_1^0 r_2^0} \left[1 + \frac{h^2}{2} \left(\frac{[\mathbf{n}_q \mathbf{v}_1]^2}{(r_1^0)^2} + \frac{[\mathbf{n}_q \mathbf{v}_2]^2}{(r_2^0)^2} + \frac{([\mathbf{n}_q \mathbf{v}_1][\mathbf{n}_q \mathbf{v}_2])}{r_1^0 r_2^0} \right) \right] \\ &+ \frac{h^2}{2} \frac{k_1^3 n_q^i [\mathbf{n}_q [\mathbf{v}_1 \mathbf{n}_q]]^j + k_2^3 n_q^j [\mathbf{n}_q [\mathbf{v}_1 \mathbf{n}_q]]^i}{q (r_1^0)^2 r_2^0} \\ &+ \frac{h^2}{2} \frac{k_1^3 n_q^i [\mathbf{n}_q [\mathbf{v}_2 \mathbf{n}_q]]^j + k_2^3 n_q^j [\mathbf{n}_q [\mathbf{v}_2 \mathbf{n}_q]]^i}{q r_1^0 (r_2^0)^2} \\ &+ \frac{h^2}{2} \frac{\delta^{ij} - n_q^i n_q^j}{q r_1^0 r_2^0}.\end{aligned}\quad (\text{A.51})$$

With degree of accuracy sufficient for our purposes,

$$\mathcal{B}_{12}^{ij} = \frac{n_q^i k_1^3 n_q^j k_2^3 c_{2,0}}{q r_1^0 (r_2^0)^2}, \quad \mathcal{C}_{12}^{ij} = \frac{n_q^i k_1^3 n_q^j k_2^3 c_{1,0}}{q (r_1^0)^2 r_2^0}. \quad (\text{A.52})$$

The calculation is straightforward, although it involves a fair amount of algebra. Finally, we obtain

$$\begin{aligned}\mathcal{B}_{12}^{ij} - \frac{\partial \mathcal{D}_{12}^{ij}}{\partial t_2} &= k_1^3 n_q^i \left(\mathcal{B}_2^j - \frac{\partial \mathcal{D}_2^j}{\partial t_2} \right) + k_2^3 n_q^j \left(\mathcal{B}_1^i - \frac{\partial \mathcal{D}_1^i}{\partial t_2} \right) \\ &- n_q^i k_1^3 n_q^j k_2^3 \left(\mathcal{B}^0 - \frac{\partial \mathcal{D}^0}{\partial t_2} \right) - k_1^3 \frac{\delta^{ij} - n_q^i n_q^j}{q^2 r_1^0}; \\ \mathcal{C}_{12}^{ij} - \frac{\partial \mathcal{D}_{12}^{ij}}{\partial t_1} &= k_2^3 n_q^j \left(\mathcal{C}_1^i - \frac{\partial \mathcal{D}_1^i}{\partial t_1} \right) + k_1^3 n_q^i \left(\mathcal{C}_2^j - \frac{\partial \mathcal{D}_2^j}{\partial t_1} \right) \\ &- n_q^i k_1^3 n_q^j k_2^3 \left(\mathcal{C}^0 - \frac{\partial \mathcal{D}^0}{\partial t_1} \right) + k_2^3 \frac{\delta^{ij} - n_q^i n_q^j}{q^2 r_2^0}.\end{aligned}\quad (\text{A.53})$$

Appendix B Bound parts of mixed energy-momentum and angular momentum

In this Appendix we consider the parts of mixed energy and mixed momentum which describe unavoidable deformations of electromagnetic “clouds” of charged particles due to mutual interaction. The short-range terms will be absorbed by four-momenta of “bare” particles within the renormalization procedure. They arise from the bound part of mixed momentum,

$$\begin{aligned}\mathcal{P}_{\text{int,bnd}}^i &= \mathcal{P}_{\text{int}}^i - \mathcal{P}_{\text{int,rad}}^i \\ &= \frac{e_1 e_2}{2} \left[\hat{\Pi}_2^i \left(\frac{\partial \sigma}{\partial t_2} \right) + \hat{\Pi}^0 (v_2^i \lambda_1) - \frac{\partial}{\partial t_1} \left(v_2^i \frac{\partial \sigma}{\partial t_2} \mathcal{D}^0 \right) \right. \\ &\quad \left. + \hat{\Pi}_1^i \left(\frac{\partial \sigma}{\partial t_1} \right) + \hat{\Pi}^0 (v_1^i \lambda_2) - \frac{\partial}{\partial t_2} \left(v_1^i \frac{\partial \sigma}{\partial t_1} \mathcal{D}^0 \right) \right],\end{aligned}\tag{B.1}$$

which is equal to the difference of the total interference momentum (3.18) and the radiative component (4.11).

Our next task is to integrate $\mathcal{P}_{\text{int,bnd}}^i$ over time variables by means of expressions (2.15) and (2.16). Recall from Section 4 that it is sufficient to evaluate the arguments of time differential operators at the ends of integration intervals.




Figure 6: Boundary points of interference of spherical wave fronts S_1 and S_2 in a hyperplane Σ_t . In momentarily comoving Lorentz frame pictured in Fig. 1 the wave fronts intersect at the coordinate origin. If $h = 0$, triangle $O_1 O_2 H$ reduces to the line. Distance q between particles is equal to the difference of radii of spheres.

In Figs. 6 the boundary conditions of interference of spherical wave fronts $S_1(O_1, k_1^0)$ and $S_2(O_2, k_2^0)$ are presented. (Fig. 3 pictures the combination of waves in four-dimensional spacetime.) Distance q between their centers, O_1 and O_2 , is equal to the difference of their




Figure 7: Acausal interference in a hyperplane Σ_t . Distance q between particles placed at centers of spheres is equal to the sum of their radii.

radii, k_1^0 and k_2^0 . In the left figure $q = k_2^0 - k_1^0$ while in the right one $q = k_1^0 - k_2^0$. Inserting this into eq.(2.5) gives $k_a^3 = k_a^0$ and $k_a^3 = -k_a^0$ for the first and the second cases, respectively.

The retarded and the advanced instants label the points on particles' world lines which are connected by a null ray. Since the relative position vector q is of null length, the null vectors K_1 and K_2 are collinear: $(K_1 \cdot K_2) = 0$. Their space parts $\mathbf{K}_1$ and $\mathbf{K}_2$ are codirectional in this case.

Within acausal region wave fronts combine in quite different manner (see Fig. 7). In contrast to the pair of points $z_1(t_1)$ and $z_2[t_2^{ret}(t_1)]$, the vertices $z_1(t_1) \in \zeta_1$ and $z_2(t_2') \in \zeta_2$ are spacelike related. Indeed, the scalar product of the separation null vectors takes minimal value $-2k_1^0 k_2^0$ if their space parts $\mathbf{K}_1$ and $\mathbf{K}_2$ are opposite directed. Hence $(q \cdot q) = 4k_1^0 k_2^0$.

In Table 1 we collect the basic quantities and functions which are evaluated at the boundary points.

To integrate the expression (B.1) over time variables, we apply the scheme developed in Section 4. Usage of the rule (4.7) implies

$$\begin{aligned}
 p_{\text{bnd},21}^i &= \frac{e_1 e_2}{2} \frac{v_2^i + n_q^i}{k_2^0 [1 - (\mathbf{n}_q \mathbf{v}_2)]} \Big|_{t_1 \rightarrow -\infty}^{[t, t_2^{ret}(t)]} + \frac{e_1 e_2}{2} \frac{v_2^i - n_q^i}{k_2^0 [1 + (\mathbf{n}_q \mathbf{v}_2)]} \Big|_{t_1 \rightarrow -\infty}^{[t_1^{ret}(t), t]} \\
 &+ \frac{e_1 e_2}{2} \frac{v_2^i - n_q^i}{k_2^0 [1 - (\mathbf{n}_q \mathbf{v}_2)]} \Big|_{[t_1^{ret}(t), t]}^{[t, t_2^{ret}(t)]}.
 \end{aligned} \tag{B.2}$$

Table 1: Relative position, retarded distances and some basic functions at the ends of integration intervals

$[t_1, t_2^{ret}(t_1)], [t_1^{adv}(t_2), t_2]$	$[t_1^{ret}(t_2), t_2], [t_1, t_2^{adv}(t_1)]$	$[t_1, t_2'(t, t_1)], [t_1'(t, t_2), t_2]$
$q = k_2^0 - k_1^0, q^0 = +q$ $k_1^3 = k_1^0, k_2^3 = k_2^0$ $r_1 = k_1^0 [1 - (\mathbf{n}_q \mathbf{v}_1)]$ $r_2 = k_2^0 [1 - (\mathbf{n}_q \mathbf{v}_2)]$	$q = k_1^0 - k_2^0, q^0 = -q$ $k_1^3 = -k_1^0, k_2^3 = -k_2^0$ $r_1 = k_1^0 [1 + (\mathbf{n}_q \mathbf{v}_1)]$ $r_2 = k_2^0 [1 + (\mathbf{n}_q \mathbf{v}_2)]$	$q = k_1^0 + k_2^0$ $k_1^3 = -k_1^0, k_2^3 = k_2^0$ $r_1 = k_1^0 [1 + (\mathbf{n}_q \mathbf{v}_1)]$ $r_2 = k_2^0 [1 - (\mathbf{n}_q \mathbf{v}_2)]$
$\sigma = 0$	$\sigma = 0$	$\sigma = -2k_1^0 k_2^0$
$\frac{\partial \sigma}{\partial t_1} = q [1 - (\mathbf{n}_q \mathbf{v}_1)]$	$\frac{\partial \sigma}{\partial t_1} = -q [1 + (\mathbf{n}_q \mathbf{v}_1)]$	$\frac{\partial \sigma}{\partial t_1} = -q [1 + (\mathbf{n}_q \mathbf{v}_1)] + 2k_2^0$
$\frac{\partial \sigma}{\partial t_2} = -q [1 - (\mathbf{n}_q \mathbf{v}_2)]$	$\frac{\partial \sigma}{\partial t_2} = q [1 + (\mathbf{n}_q \mathbf{v}_2)]$	$\frac{\partial \sigma}{\partial t_2} = -q [1 - (\mathbf{n}_q \mathbf{v}_2)] + 2k_1^0$
$\lambda_1 = k_1^0 q [1 - (\mathbf{n}_q \mathbf{v}_1)]$ $\lambda_2 = -k_2^0 q [1 - (\mathbf{n}_q \mathbf{v}_2)]$	$\lambda_1 = -k_1^0 q [1 + (\mathbf{n}_q \mathbf{v}_1)]$ $\lambda_2 = k_2^0 q [1 + (\mathbf{n}_q \mathbf{v}_2)]$	$\lambda_1 = -k_1^0 q [1 + (\mathbf{n}_q \mathbf{v}_1)]$ $\lambda_2 = -k_2^0 q [1 - (\mathbf{n}_q \mathbf{v}_2)]$

If one prefer another order of differentiation (4.8) they obtain

$$\begin{aligned}
p_{\text{bnd},12}^i &= \frac{e_1 e_2}{2} \frac{v_1^i + n_q^i}{k_1^0 [1 - (\mathbf{n}_q \mathbf{v}_1)]} \Big|_{t_2 \rightarrow -\infty}^{[t, t_2^{ret}(t)]} + \frac{e_1 e_2}{2} \frac{v_1^i - n_q^i}{k_1^0 [1 + (\mathbf{n}_q \mathbf{v}_1)]} \Big|_{t_2 \rightarrow -\infty}^{[t_1^{ret}(t), t]} \\
&+ \frac{e_1 e_2}{2} \frac{v_1^i + n_q^i}{k_1^0 [1 + (\mathbf{n}_q \mathbf{v}_1)]} \Big|_{[t, t_2^{ret}(t)]}^{[t_1^{ret}(t), t]}.
\end{aligned} \tag{B.3}$$

The lower limits $t_a \rightarrow -\infty$ vanish even if the motion is finite. Final expressions depend on particles' positions and velocities referred to the moments $t_1^{ret}(t)$ and $t_2^{ret}(t)$ as well as on the laboratory time t itself,

$$\begin{aligned}
p_{\text{bnd},21}^i &= e_1 e_2 \frac{v_2^i}{k_2^0 [1 - (\mathbf{n}_q \mathbf{v}_2)]} \Big|_{[t, t_2^{ret}(t)]} + \frac{e_1 e_2}{2} \lim_{t_2 \rightarrow t} \frac{v_2^i - n_q^i}{k_2^0 [1 + (\mathbf{n}_q \mathbf{v}_2)]} \Big|_{[t_1^{ret}(t_2), t_2]} \\
&- \frac{e_1 e_2}{2} \lim_{t_2 \rightarrow t} \frac{v_2^i - n_q^i}{k_2^0 [1 - (\mathbf{n}_q \mathbf{v}_2)]} \Big|_{[t_1'(t, t_2), t_2]};
\end{aligned} \tag{B.4}$$

$$\begin{aligned}
p_{\text{bnd},12}^i &= e_1 e_2 \frac{v_1^i}{k_1^0 [1 + (\mathbf{n}_q \mathbf{v}_1)]} \Big|_{[t_1^{ret}(t), t]} + \frac{e_1 e_2}{2} \lim_{t_1 \rightarrow t} \frac{v_1^i + n_q^i}{k_1^0 [1 - (\mathbf{n}_q \mathbf{v}_1)]} \Big|_{[t_1, t_2^{ret}(t_1)]} \\
&- \frac{e_1 e_2}{2} \lim_{t_1 \rightarrow t} \frac{v_1^i + n_q^i}{k_1^0 [1 + (\mathbf{n}_q \mathbf{v}_1)]} \Big|_{[t_1, t_2'(t, t_2)]}.
\end{aligned} \tag{B.5}$$

In an analogous way we find short-range contribution to the mixed energy due to time integration of the following expression:

$$\begin{aligned}
\mathcal{P}_{\text{int},\text{bnd}}^0 &= \mathcal{P}_{\text{int}}^0 - \mathcal{P}_{\text{int},\text{rad}}^0 \\
&= e_1 e_2 \left[\hat{\Pi}^0 \left(k_1^0 \frac{\partial \sigma}{\partial t_1} + k_2^0 \frac{\partial \sigma}{\partial t_2} + \sigma - \frac{1}{2} \frac{\partial \sigma}{\partial t_1} \frac{\partial \sigma}{\partial t_2} \right) \right]
\end{aligned} \tag{B.6}$$

$$- \frac{1}{2} \frac{\partial}{\partial t_1} \left(\frac{1}{q \| \mathbf{r}_1 \|} \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) - \frac{1}{2} \frac{\partial}{\partial t_2} \left(\frac{1}{q \| \mathbf{r}_2 \|} \frac{\partial^2 \sigma}{\partial t_1 \partial t_2} \right) \Bigg].$$

The calculation is virtually identical to that presented above, and we shall not bother with details. Finally, we obtain

$$p_{\text{bnd},21}^0 = \frac{e_1 e_2}{k_2^0 [1 - (\mathbf{n}_q \mathbf{v}_2)]} \Bigg|^{[t, t_2^{\text{ret}}(t)]} + \lim_{t_2 \rightarrow t} \frac{e_1 e_2}{k_2^0} \left[\frac{1}{1 + (\mathbf{n}_q \mathbf{v}_2)} - 1 \right]^{[t_1^{\text{ret}}(t_2), t_2]}, \quad (\text{B.7})$$

$$p_{\text{bnd},12}^0 = \frac{e_1 e_2}{k_1^0 [1 + (\mathbf{n}_q \mathbf{v}_1)]} \Bigg|^{[t_1^{\text{ret}}(t), t]} + \lim_{t_1 \rightarrow t} \frac{e_1 e_2}{k_1^0} \left[\frac{1}{1 - (\mathbf{n}_q \mathbf{v}_1)} - 1 \right]^{[t_1, t_2^{\text{ret}}(t_1)]}. \quad (\text{B.8})$$

As could be expected for bound terms, they (i) depend on the momentary state of particles' motion, (ii) contain divergent terms, and (iii) are non-covariant.

The bound components of angular momentum have similar structure,

$$M_{\text{bnd},21}^{0i} = e_1 e_2 \left[t \frac{v_2^i - n_q^i}{k_2^0 [1 - (\mathbf{n}_q \mathbf{v}_2)]} - \frac{z_2^i}{k_2^0} + \frac{t_2 n_q^i - z_2^i n_q^0}{k_2^0 [1 - (\mathbf{n}_q \mathbf{v}_2)]} \right]^{[t, t_2^{\text{ret}}(t)]} \quad (\text{B.9})$$

$$\begin{aligned} & + \frac{e_1 e_2}{2} \lim_{t_2 \rightarrow t} \left[t \frac{v_2^i + n_q^i}{k_2^0 [1 + (\mathbf{n}_q \mathbf{v}_2)]} - \frac{z_2^i}{k_2^0} - 2 \frac{t_2 n_q^i - z_2^i n_q^0}{k_2^0 [1 + (\mathbf{n}_q \mathbf{v}_2)]} \right]^{[t_1^{\text{ret}}(t), t]} \\ & - \frac{e_1 e_2}{2} \lim_{t_2 \rightarrow t} \left[t \frac{v_2^i - n_q^i}{k_2^0 [1 - (\mathbf{n}_q \mathbf{v}_2)]} - \frac{z_2^i}{k_2^0} \right]^{[t'_1(t, t_2), t_2]}, \\ M_{\text{bnd},21}^{ij} & = e_1 e_2 \frac{z_1^i v_2^j - z_1^j v_2^i}{k_2^0 [1 - (\mathbf{n}_q \mathbf{v}_2)]} \Bigg|^{[t, t_2^{\text{ret}}(t)]} \quad (\text{B.10}) \\ & - \frac{e_1 e_2}{2} (n_q^i v_2^j - n_q^j v_2^i) \left(\frac{1}{1 + (\mathbf{n}_q \mathbf{v}_2)} + \frac{1}{1 - (\mathbf{n}_q \mathbf{v}_2)} \right) \Bigg|^{[t_1^{\text{ret}}(t), t]} \\ & + \frac{e_1 e_2}{2} \lim_{t_2 \rightarrow t} \frac{z_2^i (v_2^j - n_q^j) - z_2^j (v_2^i - n_q^i)}{k_2^0 [1 + (\mathbf{n}_q \mathbf{v}_2)]} \Bigg|^{[t_1^{\text{ret}}(t_2), t_2]} \\ & - \frac{e_1 e_2}{2} \lim_{t_2 \rightarrow t} \frac{z_2^i (v_2^j - n_q^j) - z_2^j (v_2^i - n_q^i)}{k_2^0 [1 - (\mathbf{n}_q \mathbf{v}_2)]} \Bigg|^{[t'_1(t, t_2), t_2]}. \end{aligned}$$

Alternative expressions, $M_{\text{bnd},21}^{\mu\nu}$, can be obtained via reciprocity of indices 1 and 2.

In contrast to one-particle case, expanding of the expressions under limit signs in powers of $\Delta_a = t - t_a$ does not yield simple and manifestly covariant terms of clear physical sense. The “deformation” is due to the choice of the coordinate-dependent hole around the particle in the integration surface Σ_t . We neglect these structureless terms.

References

- [1] P. A. M. Dirac, Proc. R. Soc. London, Ser. A **167**, 148 (1938).

- [2] F. Rohrlich, *Classical Charged Particles* (Addison-Wesley, Redwood, CA, 1990).
- [3] C. Teitelboim, Phys. Rev. D **1**, 1572 (1970).
- [4] H. J. Bhabha, Proc. R. Soc. London, Ser. A **172**, 384 (1939).
- [5] C. A. López and D. Villarroel, Phys. Rev. D **11**, 2724 (1975).
- [6] J. M. Aguirregabiria and L. Bel, Phys. Rev. D **29**, 1099 (1984).
- [7] R. Lapiedra and A. Molina, J. Math. Phys. **20**, 1308 (1979).
- [8] R. Lapiedra, F. Marqués and A. Molina, J. Math. Phys. **20**, 1316 (1979).
- [9] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields*, 2nd ed. (Pergamon, Oxford, 1962).
- [10] J. M. Aguirregabiria and J. R. Etxebarria, J. Math. Phys. **29**, 1832 (1988).
- [11] N. P. Klepikov, Phys. Atomic Nuclei **58**, 1227 (1995); Yad.Fiz. **58**, 1307 (1995) (*in Russian*).
- [12] R. Rivera and D. Villarroel, J. Math. Phys. **38**, 5690 (1997).
- [13] R. Rivera and D. Villarroel, J. Math. Phys. **43**, 5026 (2002).
- [14] V. Hnizdo, Phys. Lett. A **129**, 426 (1988).
- [15] V. Hnizdo, J. Math. Phys. **39**, 5663 (1998).
- [16] R. Rivera and D. Villarroel, J. Math. Phys. **39**, 5664 (1998).
- [17] Yu. Yaremko, J. Phys. A: Math. Gen. **37**, L531 (2004).
- [18] Yu. Yaremko, Int. J. Mod. Phys. A **20**, 129 (2005).
- [19] B. P. Kosyakov, Phys. Rev. D **57**, 5032 (1998).
- [20] Yu. Yaremko, J. Phys. A: Math. Gen. **36**, 5149 (2003).
- [21] Yu. Yaremko, J. Phys. A: Math. Gen. **37**, 1079 (2004).
- [22] J.A. Wheeler and R.P. Feynman, Rev. Mod. Phys. **17**, 157 (1945).
- [23] J.A. Wheeler and R.P. Feynman, Rev. Mod. Phys. **21**, 425 (1949).
- [24] F. Hoyle and J.V. Narlikar, Rev. Mod. Phys. **67**, 113 (1995).
- [25] H. Spohn, Europhys. Lett. **50**, 287 (2000).
- [26] F. Rohrlich, Phys. Lett. A **283**, 276 (2001).
- [27] G. Ares de Parga, Found. Phys. **36**, 1474 (2006).

- [28] A. Ori and E. Rosenthal, Phys. Rev. D **68**, 041701(R) (2003).
- [29] A. Ori and E. Rosenthal, J. Math. Phys. **44**, 2347 (2004).